

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number
WO 02/072782 A2

(51) International Patent Classification⁷: **C12N**

(US). **TENA, Guillaume** [FR/US]; 45B Irving Street, Boston, MA 02114 (US).

(21) International Application Number: PCT/US02/07650

(74) **Agent: DECAMP, James, D.**; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110-2106 (US).

(22) International Filing Date: 12 March 2002 (12.03.2002)

(25) Filing Language: English

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) **Priority Data:**
60/275,199 12 March 2001 (12.03.2001) US

(63) **Related by continuation (CON) or continuation-in-part (CIP) to earlier application:**
US 60/275,199 (CIP)
Filed on 12 March 2001 (12.03.2001)

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) **Applicant (for all designated States except US): THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).

Published:

— without international search report and to be republished upon receipt of that report

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only): SHEEN, Jen** [US/US]; 9 Hawthorne Place, Apartment 5J, Boston, MA 02114 (US). **AUSUBEL, Frederick, M.** [US/US]; 271 Lake Avenue, Newton, MA 02161 (US). **ASAI, Tsuneaki** [JP/US]; 17 Grove Street, Apartment 5, Boston, MA 02114

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) **Title:** MASTER ACTIVATORS OF PATHOGEN RESPONSIVE GENES

(57) **Abstract:** Disclosed is a complete plant MAP kinase cascade (e.g., MEKK1, MKK4/MKK5, and MPK3/MPK6) and WRKY22/WRKY29 transcription factors that function downstream of the flagellin receptor FLS2, a leucine-rich-repeat (LRR) receptor kinase. Activation of such a MAPK cascade confers resistance to both bacterial and fungal pathogens. Also disclosed are disease-resistant plants expressing one or more members of this MAP kinase signaling cascade. Such members include constitutively active MEKK1 (Δ MEKK1), MKK4 (MKK4a), and MKK5 (MKK5a) or wild-type WRKY29.

WO 02/072782 A2

2
2

MASTER ACTIVATORS OF PATHOGEN RESPONSIVE GENES

5

Background of the Invention

The invention relates to plant disease resistance.

Despite recent progress in understanding the genetic control of plant resistance to pathogens, little progress has been reported in the identification and analysis of key regulators of pathogen resistance. Characterization of such genes would allow for the genetic engineering of plants with a variety of desirable traits. The present invention addresses these and other needs.

Summary of the Invention

15 In one aspect, the invention features a method of enhancing resistance to a plant pathogen in a plant, the method including the steps of: (a) providing a plant cell that expresses an isolated nucleic acid molecule (e.g., a DNA molecule) encoding a kinase domain of a MAPKK polypeptide; and (b) regenerating a plant from the plant cell wherein the isolated nucleic acid molecule is expressed in the plant, and wherein the plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant. In preferred embodiments, the plant is a dicot (e.g., a crucifer such as *Arabidopsis*) or a monocot. In other preferred embodiments, the kinase domain is constitutively active. And in still other embodiments, the MAPKK polypeptide is MKK4 or MKK5. Preferably, the
20 MAPKK polypeptide activates a gene involved in a pathogen defense response (for example, PAL1, GST1, WRKY29, or PR1 genes).

In another aspect, the invention features a method of enhancing resistance to a plant pathogen in a plant, the method including the steps of: (a) providing a plant cell that expresses an isolated nucleic acid molecule (e.g., a DNA molecule)
30 encoding a kinase domain of a MAPKKK polypeptide; and (b) regenerating a plant from the plant cell wherein the isolated nucleic acid molecule is expressed in the plant, and wherein the plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant. In preferred embodiments, the plant is a dicot (e.g., a crucifer such as *Arabidopsis*) or a monocot. In preferred embodiments,

the kinase domain is constitutively active, for example, a truncated form of an MAPKKK polypeptide. Preferably, the MAPKKK polypeptide is MEKK1 or ANP1. In other preferred embodiments, the MAPKKK polypeptide activates a gene involved in a pathogen defense response (for example, genes such as the PAL1, GST1, WRKY29, or PR1).

In yet another aspect, the invention features a method of enhancing resistance to a plant pathogen in a plant, the method including the steps of: (a) providing a plant cell that expresses an isolated nucleic acid molecule (e.g., a DNA molecule) encoding a polypeptide having substantial identity to a WRKY polypeptide; and (b) regenerating a plant from the plant cell wherein the isolated nucleic acid molecule is expressed in the plant, and wherein the plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant. In preferred embodiments, the plant is a dicot (e.g., a crucifer such as *Arabidopsis*) or a monocot. In preferred embodiments, the WRKY polypeptide (such as a WRKY29 or WRKY22 polypeptide) induces its own gene expression.

In another aspect, the invention features an isolated nucleic acid molecule having a nucleotide sequence for a gene promoter that initiates pathogen-inducible transcription in a plant cell, wherein the nucleotide sequence is selected from the group consisting of: (a) a nucleotide sequence including the sequence set forth in Figures 15 or 16; (b) a nucleotide sequence including at least 40 contiguous nucleotides of the sequence set forth in Figures 15 or 16; and (c) a nucleotide sequence that has at least about 70% sequence identity to a sequence set forth in (a) or (b).

In another aspect, the invention features a method for expressing a heterologous nucleotide sequence in a plant, the method including transforming a plant cell with a DNA construct including the heterologous nucleotide sequence operably linked to a promoter that is capable of initiating transcription in a plant cell and regenerating a stably transformed plant from the plant cell, wherein the promoter includes a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence including the sequence set forth in Figures 15 or 16; (b) a nucleotide sequence including at least 40 contiguous nucleotides of the sequence set forth in Figures 15 or 16; and (c) a nucleotide sequence that has at least about 70% sequence identity to a sequence set forth in (a) or (b).

In general, the kinase domain used in the methods or plants (e.g., transgenic plants or plants that are bred using a transgenic plant) of the invention is generally expressed by itself, as a MAPKKK polypeptide or a MAPKK polypeptide or kinase domain-containing fragment thereof, or as part of a genetically engineered chimeric polypeptide. Useful kinase domains include those that are capable of activating a gene involved in a plant defense response. Exemplary kinase domains include, without limitation, those that are substantially identical to the kinase domains of NPK1 or an ANP (e.g., ANP1, ANP2, or ANP3) or AtMEKK1 or MEKK4 or MEKK5.

Preferably, the methods and plants of the invention specifically utilize the kinase domain of NPK1 or ANP1. In other preferred embodiments, a full-length MAPKKK polypeptide or a kinase domain-containing fragment thereof that is substantially identical to any one of NPK1, ANP1, ANP2, or ANP3 is utilized.

The DNA encoding the constitutively active kinase domain is, in general, constitutively expressed, as is described herein. However, if desired, the kinase domain is inducibly expressed, or such a domain is expressed in a cell-specific, tissue-specific, or organ-specific manner.

In addition, WRKY polypeptides (such as WRKY29 and WRKY22) are expressed to activate plant pathogen defense responses.

Exemplary plants which are useful in the methods of the invention, as well as for generating the plants (or plant cells, plant components, plant tissues, or plant organs) of the invention, include dicots and monocots, such as sugar cane, wheat, rice, maize, sugar beet, barley, manioc, crucifer, mustard, potato, soybean, sorghum, cassava, banana, grape, oats, tomato, millet, coconut, orange, rye, cabbage, apple, grape, eggplant, watermelon, canola, cotton, carrot, garlic, onion, pepper, strawberry, yam, papaya, peanut, onion, legume, bean, pea, mango, and sunflower.

By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification (for example, glycosylation or phosphorylation).

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 40%, preferably 50%, more preferably 80%, and most preferably 90%, or even 95% sequence identity to a reference sequence (for example, the amino acid sequences of the kinase domains or full-length MAPKKK polypeptides of NPK1, MEKK1, ANP1, ANP2, or ANP3 or to their respective nucleic acid sequences, a

MAPKK polypeptides of MKK4 and MKK5 or to their respective nucleic acid sequences, or a to a WRKY polypeptide such as WRKY29 or WRKY22 or to their respective nucleic acid sequences).

For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids or greater. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides or greater.

Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705, BLAST, FastA, or PILEUP/PRETTYBOX programs). Such software matches identical or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

By "obtained from" is meant isolated from or having the sequence of a naturally-occurring sequence (e.g., a cDNA, genomic DNA, synthetic DNA, or combination thereof).

By "isolated nucleic acid molecule" is meant a nucleic acid molecule (e.g., a DNA molecule) that, is free of the genes which, in the naturally-occurring genome of the organism from which the nucleic acid of the invention is derived or obtained from, flank the gene. The term therefore includes, for example, a gene or fragment thereof that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (for example, a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a nucleic acid which is part of a hybrid gene encoding additional polypeptide sequence.

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA

molecule encoding (as used herein) a MAPKKK kinase domain (e.g., NPK1, MEKK1, ANP1, ANP2, or ANP3); a MAPKK polypeptide (e.g., MKK4 and MKK5); or a WRKY polypeptide (e.g., WRKY29 or WRKY22).

By "reporter gene" is meant a gene whose expression may be assayed; such genes include, without limitation, β -glucuronidase (GUS), luciferase (LUC), chloramphenicol transacetylase (CAT), green fluorescent protein (GFP), and β -galactosidase.

By "a promoter functional in a plant cell" is meant any minimal sequence sufficient to direct transcription in a plant cell. Included in the invention are promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell-, tissue-, or organ-specific gene expression, or elements that are inducible by external signals or agents (for example, light-, pathogen-, wound-, stress-, or hormone-inducible elements or chemical inducers) or elements that are capable of cycling gene transcription; such elements may be located in the 5' or 3' regions of the native gene or engineered into a transgene construct.

By "operably linked" is meant that a gene and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (for example, transcriptional activator proteins) are bound to the regulatory sequence(s).

By "plant cell" is meant any self-propagating cell bounded by a semi-permeable membrane and containing a plastid. Such a cell also requires a cell wall if further propagation is desired. Plant cell, as used herein, includes, without limitation, algae, cyanobacteria, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores.

By "plant component" is meant a part, segment, or organ obtained from an intact plant or plant cell. Exemplary plant components include, without limitation, somatic embryos, leaves, stems, roots, flowers, tendrils, fruits, scions, and rootstocks.

By "crucifer" is meant any plant that is classified within the Cruciferae family. The Cruciferae include many agricultural crops, including, without limitation, rape (for example, *Brassica campestris* and *Brassica napus*), broccoli,

cabbage, brussel sprouts, radish, kale, Chinese kale, kohlrabi, cauliflower, turnip, rutabaga, mustard, horseradish, and *Arabidopsis*.

By "transgene" is meant any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell which includes a nucleic acid sequence (e.g., a recombinant DNA sequence) which is inserted by artifice into a cell and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by artifice into the nuclear or plastidic genome.

By "pathogen" is meant an organism whose infection of viable plant tissue elicits a disease response in the plant tissue. Such pathogens include, without limitation, bacteria, mycoplasmas, fungi, oomycetes, insects, nematodes, viruses, and viroids. Examples of such plant diseases caused by these pathogens are described in Chapters 11-16 of Agrios, Plant Pathology, 3rd ed., Academic Press, Inc., New York, 1988.

Examples of bacterial pathogens include, without limitation, *Erwinia* (for example, *E. carotovora*), *Pseudomonas* (for example, *P. syringae*), and *Xanthomonas* (for example, *X. campestris* and *X. oryzae*).

Examples of fungal or fungal-like disease-causing pathogens include, without limitation, *Alternaria* (for example, *A. brassicola* and *A. solani*), *Ascochyta* (for example, *A. pisi*), *Botrytis* (for example, *B. cinerea*), *Cercospora* (for example, *C. kikuchii* and *C. zaea-maydis*), *Colletotrichum* sp. (for example, *C. lindemuthianum*), *Diplodia* (for example, *D. maydis*), *Erysiphe* (for example, *E. graminis* f.sp. *graminis* and *E. graminis* f.sp. *hordei*), *Fusarium* (for example, *F. nivale* and *F. oxysporum*, *F. graminearum*, *F. solani*, *F. moniliforme*, and *F. roseum*), *Gaeumanomyces* (for example, *G. graminis* f.sp. *tritici*), *Helminthosporium* (for example, *H. turcicum*, *H. carbonum*, and *H. maydis*), *Macrophomina* (for example, *M. phaseolina* and *Maganaporthe grisea*), *Nectria* (for example, *N. heamatocacca*), *Peronospora* (for example, *P. manshurica*, *P. tabacina*), *Phoma* (for example, *P. betae*), *Phymatotrichum* (for example, *P. omnivorum*),

Phytophthora (for example, *P. cinnamomi*, *P. cactorum*, *P. phaseoli*, *P. parasitica*,
P. citrophthora, *P. megasperma f.sp. sojae*, and *P. infestans*), *Plasmopara* (for
example, *P. viticola*), *Podosphaera* (for example, *P. leucotricha*), *Puccinia* (for
example, *P. sorghi*, *P. striiformis*, *P. graminis f.sp. tritici*, *P. asparagi*, *P. recondita*,
5 and *P. arachidis*), *Puthium* (for example, *P. aphanidermatum*), *Pyrenophora* (for
example, *P. tritici-repentens*), *Pyricularia* (for example, *P. oryzae*), *Pythium* (for
example, *P. ultimum*), *Rhizoctonia* (for example, *R. solani* and *R. cerealis*),
Scerotium (for example, *S. rolfsii*), *Sclerotinia* (for example, *S. sclerotiorum*),
Septoria (for example, *S. lycopersici*, *S. glycines*, *S. nodorum* and *S. tritici*),
10 *Thielaviopsis* (for example, *T. basicola*), *Uncinula* (for example, *U. necator*),
Venturia (for example, *V. inaequalis*), and *Verticillium* (for example, *V. dahliae* and
V. albo-atrum).

Examples of pathogenic nematodes include, without limitation, root-knot
nematodes (for example, *Meloidogyne sp.* such as *M. incognita*, *M. arenaria*, *M.*
15 *chitwoodi*, *M. hapla*, *M. javanica*, *M. graminicola*, *M. microtyla*, *M. graminis*, and
M. naasi), cyst nematodes (for example, *Heterodera sp.* such as *H. schachtii*, *H.*
glycines, *H. sacchari*, *H. oryzae*, *H. avenae*, *H. cajani*, *H. elachista*, *H. goettingiana*,
H. graminis, *H. mediterranea*, *H. moths*, *H. sorghi*, and *H. zaeae*, or, for example,
Globodera sp. such as *G. rostochiensis* and *G. pallida*), root-attacking nematodes
20 (for example, *Rotylenchulus reniformis*, *Tylenchulus semipenetrans*, *Pratylenchus*
brachyurus, *Radopholus citrophilus*, *Radopholus similis*, *Xiphinema americanum*,
Xiphinema rivesi, *Paratrichodorus minor*, *Heterorhabditis heliothidis*, and
Bursaphelenchus xylophilus), and above-ground nematodes (for example, *Anguina*
funesta, *Anguina tritici*, *Ditylenchus dipsaci*, *Ditylenchus myceliphagus*, and
25 *Aphenlenchoides besseyi*).

Examples of viral pathogens include, without limitation, tobacco mosaic
virus (TMV), tobacco necrosis virus (TNV), potato leaf roll virus, potato virus X,
potato virus Y, tomato spotted wilt virus, and tomato ring spot virus.

By "enhanced resistance to a plant pathogen" is meant a level of resistance to
30 a disease-causing pathogen in a non-naturally occurring plant (or cell or seed
thereof) which is greater than the level of resistance in a control plant (for example,
a non-transgenic plant or wild-type). In preferred embodiments, the level of
resistance in a non-naturally occurring transgenic plant of the invention is at least

5% to 20% (and preferably 30% or 40%) greater than the resistance exhibited by a control plant. In other preferred embodiments, the level of resistance to a disease-causing pathogen is 50% greater, 60% greater, and more preferably even 75% or 90% greater than a control plant; with up to 100% or greater above the level of resistance as compared to a control plant being most preferred. The level of resistance is measured using conventional methods. For example, the level of resistance to a pathogen may be determined by comparing physical features and characteristics (for example, plant height and weight) or by comparing disease symptoms (for example, delayed lesion development, reduced lesion size, leaf wilting and curling, water-soaked spots, amount of pathogen growth, and discoloration of cells) of the non-naturally occurring plant (e.g., a transgenic plant).

By "detectably-labeled" is meant any direct or indirect means for marking and identifying the presence of a molecule, for example, an oligonucleotide probe or primer, a gene or fragment thereof, or a cDNA molecule or a fragment thereof. Methods for detectably-labeling a molecule are well known in the art and include, without limitation, radioactive labeling (for example, with an isotope such as ³²P or ³⁵S) and nonradioactive labeling (for example, fluorescence of chemiluminescent labeling, for example, fluorescein labeling).

The invention provides for increased production efficiency, as well as for improvements in quality and yield of crop plants and ornamentals. Thus, the invention contributes to the production of high quality and high yield agricultural products; for example, fruits, ornamentals, vegetables, cereals, and field crops. Genetically-improved seeds and other plant products that are produced using plants expressing the genes and methods described herein also render farming possible in areas previously unsuitable for agricultural production. The invention further provides a means for mediating the expression of pathogen defense response genes (e.g., GST1, PAL, WRKY29, and PR1) that enable a plant to resist its pathogens. For example, transgenic plants constitutively expressing a kinase domain of a MAPKKK or MAPKK, or a WRKY polypeptide are capable of turning on a plant's pathogen defense transduction pathway by activating the expression of plant pathogen defense regulatory pathways.

As discussed above, MAPK activation is responsible for providing plants with the ability to protect themselves against pathogens have been identified.

Accordingly, the invention provides a number of important advances and advantages for the protection of plants against their pathogens. For example, expression of such genes *in planta*, as described herein, facilitates an effective and economical means for in-plant protection against plant pathogens. Such protection against pathogens
5 reduces or minimizes the need for traditional chemical practices (for example, application of fungicides, bactericides, nematocides, insecticides, or viricides) that are typically used by farmers for controlling the spread of plant pathogens and providing protection against disease-causing pathogens.

The invention further provides a means for mediating the expression of
10 pathogenesis-related proteins, for example, GST, that confers resistance to plant pathogens. For example, transgenic plants constitutively producing an MAPK cascade gene product (e.g., a MAPKKK, MAPKK, or WRKY polypeptide) are capable of activating PR gene expression, which in turn confers resistance to plant pathogens. Collective PR gene expression that is mediated by expression of a
15 MAPK cascade polypeptide (e.g., a MAPKKK, MAPKK, or WRKY polypeptide) obviates the need to express individual resistance genes as a means to promote plant defense mechanisms.

Moreover, constitutively active derivatives of MAPKKKs (e.g., MEKK1 and ANP1) and MAPKKs (e.g., MKK4 and MKK5) can activate the *PAL1*, *GST1*, and
20 *WRKY29* and *PR1* gene promoters. Thus, MEKK1, ANP1, MKK4 and MKK5 (and their orthologs in other plants) activate the same MAPK cascades. These PKs are therefore reasonable targets for transgenic plant manipulation to generate agronomically valuable traits, such as broad-spectrum pathogen resistance. The *PAL1* gene encodes phenylalanine ammonia lyase that functions at a universal key
25 step for the biosynthesis of salicyclic acid and phytoalexins (anti-pathogen hormones and chemicals). The expression of GST1 (glutathione S-transferase) is important for the protection and detoxification of cells during pathogen infection. The activation of the WRKY29 (a transcription activator and DNA binding protein for the W box) provides a positive feedback loop for the regulation of *PAL1* and
30 *GST1* genes. The PR1 protein has an important anti-pathogen activity.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

The drawings will first be described.

Figure 1 shows a MAPK signaling pathway leading to WRKY29.

Figure 2 shows a schematic diagram of the *Arabidopsis* WRKY29 polypeptide, including the WRKY domain and its W box target sequence.

Figure 3 shows that Flg22 induces WRKY29 in *Arabidopsis* protoplasts.

Figure 4 shows the MKK4 cDNA sequence and its predicted amino acid sequence.

Figure 5 shows the strategy used to render MKK4 constitutively active.

Figure 6 shows the MKK5 cDNA sequence and its predicted amino acid sequence.

Figure 7 shows the strategy used to render MKK5 constitutively active.

Figure 8 shows the alignment of MKK4 and MKK5 polypeptides.

Figure 9 shows that constitutively active AtMEKK1 induces WRKY29.

Figure 10 shows that constitutively active AtMKK4 induces WRKY29.

Figure 11 shows that WRKY29 induces its own promoter.

Figure 12 shows that WRKY29 regulates early defense genes. Pal1 refers to phenylalanine ammonia lyase; FRK1 refers to the flg22 induced receptor-like kinase; GST1 refers to glutathione S-transferase 1; and GST6 refers to glutathione S-transferase 6.

Figure 13 shows a schematic diagram of the *Agrobacterium*-mediated transient transfection protocol.

Figure 14 shows that transient expression of WRKY29 reduces susceptibility to *Pseudomonas syringae*.

Figure 15 shows the pathogen-inducible WRKY29 gene promoter fragment.

Figure 16 shows the pathogen-inducible WRKY22 gene promoter fragment.

Figures 17A-D show the early defense gene activation by flg22. Figure 17A shows a RT-PCR analysis. Protoplasts were preincubated for one hour with (+) or without (-) 10 μ M CHX, and then treated without or with 100 nM flg22. Figure 17B shows the involvement of MAPK. Transfected protoplasts were preincubated for one hour with 1 μ M staurosporine (ST) or with 1 μ M U0126 before induction by 100 nM flg22 for four hours. Figure 17C shows the FLS2 requirement. Protoplasts isolated from wild-type (*F*) and *fls2-24* mutant (*f*) leaves were transfected with an

effector plasmid expressing GFP (-) or FLS2. Transfected protoplasts were incubated for nine hours first before incubation with or without 100 nM flg22 for four hours. Figure 17D shows MKP1 suppression. Protoplasts were transfected with an effector plasmid expressing GFP (-) or MKP1.

5 Figures 18A-D show that Flg22 activates MPK3 and MPK6 through FLS2. Figure 18A shows that Flg22 activates endogenous MBP kinases. Protoplasts were treated with 1 μ M flg22 (+) or water (-). Figure 18B shows that Flg22 activates MPK3 and MPK6. MAPK activation with or without 100 nM flg22 for 10 minutes (top). Expression of each MAPK (2, 3, 5, 6, 7 or 9) protein was verified by
10 [³⁵S]methionine labeling and immunoprecipitation (bottom). Figure 18C shows that Flg22 activation of MPK3 and MPK6 requires FLS2. A HA-tagged MAPK (3 or 6) was coexpressed with a wild-type (FLS2) or kinase-inactive (FLS2 Km) flagellin receptor with a HA tag in *fls2-24* protoplasts. MAPK activities (top) and FLS2 (middle) and MAPK (bottom) protein levels are shown. Control experiment was
15 performed with wild-type (Wt) protoplasts. Figure 18D shows that MKP1 abolishes flg22 activation of MPK3 and MPK6. MAPK activity (top), and MAPK (middle), and MKP1 (bottom) protein levels are shown.

 Figures 19A-E show the MKK4 and MKK5 activation of MPK3 and MPK6 and the flg22-inducible early defense genes. Figure 19A shows that MKK5
20 activates MPK3 and MPK6. Protoplasts were transfected with plasmids expressing MAPK (2, 3, 5, 6, 7 or 9) and wild-type MKK5 (w) or constitutively active MKK5 mutant (a). MAPK activity (top) and MAPK (middle), and MAPKK protein levels are shown (bottom). Asterisk indicates a background band. Figure 19B shows that MKK4 and MKK5 are redundant for MPK3 and MPK6 activation. Protoplasts were
25 transfected with plasmids expressing MAPK (3 or 6) and one of four MYC-tagged and constitutively active MAPKK mutants (1, 2, 4, or 5) or a control plasmid (-). MAPK activity (top), MAPK (middle) and MAPKK (bottom) protein levels are shown. Figure 19C shows that MKK4a and MKK5a activate *WRKY29* and *FRK1* promoter activity. Protoplasts were expressing GFP (-) or a constitutively active
30 MKK (1, 2, 4, or 5). Figure 19D shows that dominant-negative mutants of MKK4 and MKK5 partially inhibit flg22 activation of the *WRKY29* and *FRK1* promoters. Protoplasts were expressing GFP (-) or a kinase-inactive mutant of MKK4 or MKK5

(4 or 5). Figure 19E shows that MKK4 and MKK5 act downstream of FLS2. Wild-type (*FLS2*) and *fls2-24* (*fls2*) protoplasts were expressing GFP (-) or a constitutively active mutant of MKKs.

Figures 20A-D show that MEKK1 initiates the flg22 MAPK cascade. Figure 5 20A shows that constitutively active MEKK1 activates MKK5. Protoplasts were expressing wild-type MKK5 or a control plasmid (-), and one of four HA-tagged constitutively active MAPKKs (kinase domain) or a control plasmid (-). MKK5 activity based on GST-MPK6Km as a substrate (top) and MAPKKK (middle) and MAPKK (bottom) protein levels are shown. Figure 20B shows that constitutively 10 active MEKK1 induces the *WRKY29* and *FRK1* promoters. Protoplasts were expressing GFP (-) or constitutively active MEKK1 (M) or CTR1 (C). Figure 20C shows that MEKK1 acts downstream of FLS2. Protoplasts were isolated from wild-type (*FLS2*) and mutant (*fls2*) leaves. Figure 20D shows that dominant-negative mutant of MEKK1 inhibits flg22 activation of the *WRKY29* and *FRK1* promoters. 15 Protoplasts were expressing a kinase inactive mutant of the full-length MEKK1 (MEKK1in).

Figures 21A-F show that the flg22 MAPK cascade and WRKY29 are important for *Arabidopsis* defense signaling. Figure 21A shows that WRKY29-GFP is localized in the nucleus. Visualization of protoplasts expressing WRKY29-GFP 20 (top), a red-fluorescent nuclear marker (middle), and super-imposition (bottom), in the absence of flg22. Figure 21B shows that WRKY22 and WRKY29 activate early defense genes. Protoplasts were expressing GFP (-), WRKY29 ("29"), WRKY22 ("22") or WRKY42 ("42"). Figure 21C shows that WRKY29 acts downstream of FLS2. Protoplasts were isolated from wild-type (*FLS2*) and mutant (*fls2*) leaves. 25 Figure 21D shows that *Arabidopsis* leaves expressing Δ MEKK1, MKK4a, MKK5a, or WRKY29 exhibited reduced disease symptoms after *P. syringae* (Ps) infection. The control leaves were infiltrated with 10 mM MgSO₄ (Mg). Figure 21E shows *Arabidopsis* leaves expressing Δ MEKK1, MKK4a, or WRKY29 exhibit reduced disease symptoms after *B. cinerea* infection. The right half of each leaf was infected 30 with *B. cinerea*. Figure 21F shows the early stage of *B. cinerea* development was inhibited on leaves expressing MKK4a. On MKK1a-expressing leaves (MKK1a), germinated spores (s) of *B. cinerea* formed superficial hyphae (h) and branched

appressoria (a) two days after infection. The fungal spores (s) formed only germ tubes (gt) on MKK4a-expressing leaves (MKK4a). Bars: 50 μ M.

Figure 22 shows a model of innate immune signaling activated by LRR receptors in *Arabidopsis*, mammals, and *Drosophila*. A putative repressor (R) could control WRKY22 and WRKY29 activity since their overexpression bypasses the requirement of elicitors. The conserved signaling pathways for innate immune responses in animals are summarized based on recent reviews on mammals and *Drosophila*.

10 Detailed Description

MAPK cascades are, in general, conserved in eukaryotes. Genome sequence comparisons and molecular analysis of signal transduction pathways have revealed the existence of conserved MAPK signaling cascades in eukaryotes that connect environmental and developmental signals to the activation of a variety of species-specific downstream genes. Extensive studies have revealed the functions of five yeast MAPKs in stress, pheromone, and nutrient responses. Based mostly on analysis in cellular systems, four distinct classes of animal MAPKs have been shown to participate in signaling pathways triggered by growth regulators, neurotransmitters, cytokines, and diverse stresses. It appears likely that additional MAPK signaling pathways will be identified. BLAST searches of DNA databases that have been conducted using a conserved signature motif that covers the phosphorylation site TXY of MAPKs identified relatively large families of putative MAPK genes in yeast (6 genes), *C. elegans* (17 genes), mouse (17 genes), human (27 genes), and *Arabidopsis* (18 genes). In the case of *Arabidopsis*, there are at least 7 putative MAPKK genes and 29 MAPKKK genes. Homologs of many of these genes can be found in other plant species. However, the functions of most of these putative MAPK cascade components have not been determined. As is detailed herein, several of these genes are actively involved in regulating the expression of pathogen defense mechanisms.

30 For cell signaling, down regulation of the MAPK activity is just as important as its up regulation. Inactivation of MAPK can be achieved through dephosphorylation of crucial protein tyrosine and threonine residues by dual-specificity MAPK phosphatases (MKPs), protein tyrosine phosphatases (PTPs), and

serine/threonine protein phosphatases PP2C. Homologs of MKP, PTP and PP2C genes have been identified in yeast, *C. elegans*, *Arabidopsis*, and mammals.

Depending on their tissue specificity, mode of induction, subcellular localization, and substrate preferences, each PP might participate in the repression of a specific

5 MAPK activity. Signals that activate MAPKs can often activate the transcription of MKPs or PTPs, that might be important for the negative feedback regulation of the MAPK pathways. The roles of most plant MKPs, PTPs, and PP2Cs in MAPK cascades are unknown.

As is detailed below, an *Arabidopsis* protoplast transient expression assay
10 and four reporter genes have been used to identify protein (PKs) that can activate the early pathogen response genes, GST1, PAL1, and WRKY29, as well as a late response gene, PR1. Constitutively active derivatives of two MAPKKKs (MEKK1 and ANP1) and two MAPKKs (MEK4 and MEK5) activated the PA11, GST1, WRKY29 and PR1 gene promoters. Accordingly, MEKK1, ANP1, MEK4 and
15 MEK5 activate the same MAPK cascades. These PKs are therefore reasonable targets for transgenic plant manipulation to generate agronomically valuable traits, such as broad-spectrum pathogen resistance. In particular, one MAPK cascade involving Flg22 peptide-mediated defense responses has been examined (Figure 1). The overall strategy involved the identification and cloning of several *Arabidopsis*
20 genes encoding MAPKs, MAPKKs, MAPKKKs, and WRKYs. Transient expression of epitope-tagged MAPKs, MAPKKs, and MAPKKKs were then used to determine which MAPK cascades respond to particular signals in *Arabidopsis* and maize protoplasts. Engineered MAPKKKs and MAPKKs designed to be constitutively active or act as dominant negative inhibitors, as well as WRKYs, were
25 used to identify specific reporter genes associated with particular signaling pathways. Using this system, we have identified a complete plant MAPK cascade and WRKY transcription factors acting downstream of the flagellin receptor FLS2. The data presented below also indicate that this signaling pathway functions in response to both bacterial and fungal pathogens and could therefore be engineered to
30 enhance the disease resistance of crop plants to a wide range of pathogens.

MAPK Activation in the Plant Defense Response

In response to pathogenic microorganisms, plants induce a variety of cellular reactions. Among the reactions are the induction of rapid programmed cell death (PCD), known as the hypersensitive response (HR), and the transcriptional
5 activation of many defense-related genes, including the early GST1 (glutathione S-transferase) and PAL1 (phenylalanine ammonia-lyase) genes, as well as the later pathogenesis-related (PR) genes. Activation of these defense responses is often governed by a specific "gene-for-gene" interaction between a plant resistance (R) gene and a pathogen avirulence (avr) gene. Despite significant insights into R gene
10 structure as putative receptors, little is known about the signal transduction events following R-Avr interactions. The finding that two R gene products, tomato PTO and rice XA-21, contain a protein kinase domain suggest that protein phosphorylation might play a central role in R gene-dependent signal transduction. More recent studies have provided compelling evidence that MAPKs are likely
15 involved in this signaling event. In tobacco, the N gene confers resistance to tobacco mosaic virus (TMV). It was demonstrated that two previously identified MAPKs, a SA-induced protein kinase (SIPK) and a wound-induced protein kinase (WIPK), were activated during infection by TMV. Importantly, WIPK activation by TMV depended on N since the activation was not detected in tobacco plants lacking
20 the N gene. Similarly, treatment of tobacco suspension cells expressing the tomato R gene, Cf-9, with the corresponding avr gene product, Avr9, resulted in rapid activation of both tobacco SIPK and WIPK in a strictly gene-for-gene manner.

Involvement of MAPKs in plant defense signaling has also been demonstrated through the analysis of plant responses after non-specific elicitation
25 with bacterial or fungal oligosaccharides, proteins, or peptides. Parsley cells recognize a fungal pathogen *Phytophthora sojae* through a plasma membrane receptor. A pathogen-derived oligopeptide elicitor binds to this receptor and thereby stimulates a multi-component defense responses. It was demonstrated that the elicitor activates a MAPK that is similar to tobacco WIPK. A regulatory function of
30 a MAPK in defense gene induction has also been suggested in tobacco suspension cells, in which three fungal elicitors rapidly activate the SIPK.

Another important virulence factor for bacteria that are pathogenic to animals and plants is the flagellum. It has been demonstrated that plants have a

highly sensitive chemoperception system for eubacterial flagellins, specifically targeted to the most highly conserved domain within its N terminus. Synthesis of this peptide (22 amino acids) creates a highly active elicitor called Flg22. In *Arabidopsis* seedlings, Flg22 elicits callose deposition, induction of PR genes, and growth inhibition. Synthetic peptides, carrying the corresponding flagellin sequences of bacteria that coexist intimately with plants such as *Rhizobium meliloti* and *Agrobacterium tumefaciens*, fail to induce any defense responses. It was recently reported that, in *Arabidopsis*, one of the loci conferring Flg22 sensitivity encodes a putative transmembrane protein kinase with leucine-rich repeats, resembling the rice resistance gene XA-21. As is demonstrated herein, the MAPK pathway is involved in Flg22 signaling. There now follow particular examples of the invention. These examples are provided for the purpose of illustrating the invention, and should not be construed as limiting.

Example 1: Cellular Systems for Functional Analysis of Plant Defense Gene Regulatory Pathways

Two protoplast transient expression systems, such as the maize (Sheen, 1999) or *Arabidopsis* (Kovtun et al, *Proc. Natl. Acad. Sci., USA* 97:2940-5, 2000) leaf protoplast systems, are useful for examining regulation of plant defense mechanisms. Technical advances in these two systems, including high transformation efficiency by electroporation (up to 75%) or by PEG fusion (up to 80%), specificity of reporter gene regulation, the use of improved green-fluorescent protein (GFP) as a vital and visual reporter (Chiu et al., *Curr. Biol.* 6, 325-30, 1996), PK and PP activity assays (Kovtun et al., *Nature* 395:716-720, 1998; Sheen, *Proc. Natl. Acad. Sci., USA* 95:975-980, 1998; Kovtun et al., *Proc. Natl. Acad. Sci., USA* 97:2940-5, 2000), and expression of putative signaling molecules with epitope- or GFP-tag and their detection by immunoprecipitation and confocal microscopy, make them especially attractive to examine MAPK cascades and the plant defense gene regulatory pathways.

Because MAPK cascades are regulators of pathogen defense the manipulation of such MAPK cascades in plants provides new tools for crop improvement in disease resistance. Furthermore, the high degree of conservation of MAPK signaling pathways facilitates the transfer of knowledge learned in the *Arabidopsis* system to agricultural species.

Example 2: Flg22-induced Responses in *Arabidopsis* Protoplasts

Activation or repression of the transcription of specific reporter genes can be used to monitor activation of particular signaling pathways. For example, Flg22 has been found to induce the expression of the GST1, PAL1, and PAL2 promoters.

5 MAPK activation was demonstrated in *Arabidopsis* protoplasts isolated from 4-week-old leaves. Here Flg22 or distilled water was added to the protoplasts and samples were assayed at different times to determine relative promoter activity. Defense gene promoters fused to a luciferase reporter gene were introduced into *Arabidopsis* protoplasts. The protoplasts were incubated for 16 hours in the

10 presence of Flg22 and/or staurosporine, and luciferase activities were assayed. The number of viable cells in each sample was also determined at the end of the incubation by Evans blue staining and promoter activity was represented as a luciferase activity per viable cell. In the Flg22 experiment, the induction of all the promoters was effectively suppressed by staurosporine, a protein kinase inhibitor.

15 However, the GST6 promoter was not induced by Flg22.

Example 3: Flg22 induces WRKY29 in *Arabidopsis* protoplasts and a MAPK Signal Cascade

Arabidopsis protoplasts treated Flg22 were evaluated for the expression of WRKY29 using standard reverse-transcription polymerase chain reaction analysis

20 (RT-PCR). As shown in Figure 3, Flg22 was found to induce WRKY29. The structure of WRKY29 is shown in Figure 2. To determine whether, MAP kinase signaling is involved in WRKY29 induction, the induction of WRKY29 was monitored in the presence of U0126, a MAPKK inhibitor. RT-PCR analysis of WRKY29 induction showed that U0126 suppressed WRKY29 induction.

25 Furthermore, Flg22 was found to activate MBP kinases. Further demonstrating the involvement of a MAPK cascade, mouse MAPK phosphatase1 was found to suppress WRKY induction by Flg22.

Example 4: Constitutively Active ANP1 and MEKK1 Activate PAL1, GST1, and WRKY29

30 To determine whether ANP1 and MEKK1 were involved in activating plant defense response genes, the effect of a constitutively active ANP1 and MEKK1 on the activity of the PAL1 promoter was examined. These polypeptides were rendered constitutively active according to standard methods known in the art. It has been shown that MAPKKKs consist of a well-conserved kinase domain and putative

regulatory domains. Truncated or naturally occurring MAPKKs carrying only the kinase domain have been shown to have constitutive kinase activity. The expression of the constitutively active ANP1 and MEKK1 was found to activate PAL1 gene expression. In addition, constitutively active ANP1 and MEKK1 was found to
5 activate the GST1 and WRKY29 gene promoters (Figure 9).

Example 5: Constitutively Active AtMKK4 and AtMKK5 Activate PAL1 and WRKY29

To determine whether MKK4 and MKK5 are involved in activating plant defense response genes, the effect of a constitutively active AtMKK4 and MKK5 on
10 the activity of the WRKY29 promoter was examined. The nucleotide and predicted amino acid sequences of MKK4 and MKK5 are shown in Fig 4 and 6, respectively. The strategy for rendering each of these PKs constitutively active is also shown
Figures 5 and 7. Furthermore, an alignment of MKK4 and MKK5 is shown in Figure 8. The expression of constitutively active MKK4 and MKK5 was found to
15 activate PAL1 and WRKY29 gene expression (Figure 10).

Example 6: WRKY29 Induces Its Own Promoter

To determine the role of WRKY29 in plant defense, the effect of expressing WRKY29 on the activation of a reporter gene under the transcriptional control of the WRKY29 promoter was examined. As shown in Figure 11, the WRKY29
20 polypeptide was found to induce expression of its own gene promoter. Exemplary pathogen-inducible WRKY promoter fragments are shown in Figures 15 and 16. In addition, as shown in Figure 12, WRKY29 was found to regulate PAL1 and the so-called flg22 induced receptor-like kinase, but not a GST gene promoter.

Example 7: Transient Expression of WRKY29 Reduces Susceptibility to Pseudomonas syringae

To determine whether expression of WRKY29 reduces pathogen susceptibility, *Arabidopsis* leaves from 4-5 week-old plants were infiltrated with *Agrobacteria* carrying 35S-GFP, 35S-MKK4act, or 35S-WRKY29 (Figure 13). The plants were incubated for three days and the *Agrobacterium*-infiltrated leaves were
30 further infiltrated with *Pseudomonas syringae* pv. *maculicola* strain ES4326 (resuspended in 10 mM MgSO₄, OD₆₀₀ = 0.002) or 10 mM MgSO₄. The plants were incubated for two more days and the infiltrated leaves were cut for examination. As shown in Figure 14, transient expression of WRKY29 was found to reduce susceptibility to *Pseudomonas syringae* pv. *maculicola* strain ES4326.

Using the same procedure, it was also found that transient expression of a constitutively active MKK4 reduced the plant's susceptibility to *Pseudomonas syringae*.

5 Example 8: Early defense gene transcription induced by flg22 in *Arabidopsis* leaf cells

To dissect the early signal transduction pathways in plant innate immune responses, we first established a leaf cell assay based on flg22 inducible transcription of early response genes in *Arabidopsis* mesophyll protoplasts. Few reporter genes have been developed for the dissection of the early stages of defense signaling pathways in plants (Durrant et al., *Plant Cell* 12:963-977, 2000; Maleck et al., *Nature Genet.* 26:403-410, 2000; Schenk et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:11655-11660, 2000). To identify genes that are induced by flg22-activated defense signaling, we created a subtracted cDNA library that represented mRNA species induced at various times by the elicitor flg22 in *Arabidopsis* mesophyll protoplasts according to the methods described by Asai et al. (*Plant Cell* 12:1823-1835, 2000). Using flg22 rather than a pathogen or a natural elicitor avoided the possibility that multiple elicitors could be functioning in parallel. Furthermore, synchronous elicitation by flg22 is achieved more reproducibly in homogeneous mesophyll protoplasts than in intact leaves. In the library we found well-described defense genes, such as *PAL1* (At2g37040), *GST1* (At1g02930), *PR1* (At2g19990), and *PR5* (At1g75040), which are induced by a variety of pathogens and elicitors at different stages of the defense response in many plant species (Maleck et al., *Nature Genet.* 26:403-410, 2000; Schenk et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:11655-11660, 2000; Asai et al., *Plant Cell* 12:1823-1835, 2000). These genes were also induced in *Arabidopsis* leaves infiltrated with flg22 (data not shown), suggesting that similar defense responses are induced by flg22 in isolated leaf protoplasts and in leaves of intact plants.

To develop new reporter genes that are expressed early in the primary defense response, we focused on flg22-activated genes identified in the subtraction library that were expressed at early time points (see below) and encode putative regulatory factors, including *WRKY29* (At4g23550) (Euglem et al., *Trends Plant Sci.* 5:199-206, 2000) and *FRK1* (*FLG22-INDUCED RECEPTOR-LIKE KINASE 1*, At2g19190), which encode a WRKY transcription factor (with a conserved WRKY

DNA-binding domain) and a LRR receptor kinase, respectively. Induction of WRKY transcription factors and LRR receptor kinase genes by pathogens, pathogen-derived elicitors, or salicylic acid has been demonstrated in the leaves of several plant species including *Arabidopsis* (data not shown). Taken together, these results indicate that *Arabidopsis* protoplasts treated with flg22 increase transcription of a variety of defense-related genes that are also expressed in intact plants after pathogen infection.

Polymerase chain reaction after reverse transcription of RNA (RT-PCR) analysis showed that *WRKY29*, *FRK1*, and *GST1* mRNA levels were elevated in *Arabidopsis* protoplasts within 30 minutes after flg22 treatment (Fig. 17A), whereas induction of the extensively studied *PR1* and *PR5* genes occurred much later (Fig. 17A). Two stress-regulated genes, H₂O₂-inducible *GST6* (At2g47730) and ABA-, cold-, or drought-responsive *RD29A* (AT5g52310) (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000), were not induced by flg22 over basal levels of expression (Fig. 17A). In contrast to the common assumption that protoplasts isolated from fresh leaves are badly damaged, stressed and dying, or are already activated in defense responses, the data in Figure 17 (as well as the data presented below) show that *Arabidopsis* mesophyll protoplasts respond specifically to flg22, similarly to the response observed in intact plants (Gomez-Gomez et al., *Plant J.* 18:277-284, 1999). The specific response of mesophyll protoplasts to flg22 is consistent with our previous studies using mesophyll protoplasts as a versatile system to analyze diverse stress signaling pathways (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000; Sheen et al., *Science* 274:1900-1902, 1996) and cell death programs (Asai et al., *Plant Cell* 12:1823-1835, 2000).

To examine whether *de novo* protein synthesis is required for flg22-induced defense gene expression, protoplasts were preincubated with the protein synthesis inhibitor cycloheximide (CHX) and then treated with flg22. RT-PCR analysis showed that induction of the early response genes *WRKY29*, *FRK1*, and *GST1* by flg22 was not significantly affected by CHX, whereas the inhibitor blocked induction of the late response genes *PR1* and *PR5*, but not the expression of other control genes (Fig. 17A). To determine whether the early defense genes are activated transcriptionally, the promoters of the *WRKY29*, *FRK1*, *GST1*, *GST6*, and *RD29A* genes were fused to the *LUC* reporter gene as described by Kovtun et al.

(*Proc. Natl. Acad. Sci., USA* 97:2940-2945, 2000) and tested for their response to flg22 in transiently transfected protoplasts. Consistent with the RT-PCR data obtained with the endogenous genes, the *WRKY29*, *FRK1*, and *GST1* promoters, but not the *GST6* and *RD29A* promoters, were induced by flg22 (Fig. 17B). In a parallel control experiment, the 23-amino acid peptide flg23RM, corresponding to the flg22-homologous sequence of the *Rhizobium melliloti* flagellin protein, activated none of these promoters (data not shown). *R. melliloti* is a plant symbiont that fails to activate (and/or suppress) host innate immunity. In previous work, *R. melliloti* flagellin was shown not to activate host defense responses (Felix et al., *Plant J.* 18:265-276, 1999; Gomez-Gomez et al., *Plant J.* 18:277-284, 1999). These results suggest that in response to pathogen-derived flagellin, *WRKY29*, *FRK1*, and *GST1* are rapidly induced by pre-existing signaling molecules. Because the H₂O₂-responsive gene *GST6* (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000) was not induced by flg22 (Figs. 17A and 17B), it is likely that although flg22 induces an oxidative burst in *Arabidopsis* leaves (Gomez-Gomez et al., *Mol. Cell* 5:1003-1011, 2000; Gomez-Gomez et al., *Plant J.* 18:277-284, 1999), it may not elicit a strong or persistent oxidative burst in protoplasts. Consistent with this conclusion, an inhibitor of H₂O₂ accumulation, diphenylene iodonium, had no effect on flg22-induced *WRKY29*, *FRK1* and *GST1* gene expression (data not shown). Thus, the flg22-protoplast system distinguishes between flagellin and oxidative-stress signaling pathways (see below). In addition to the well-established parsley protoplast system derived from suspension cultures (Nurnberger et al., *Trends Plant Sci.* 6:372-379, 2001; Blume et al., *Plant Cell* 12:1425-1440, 2000; Hirt et al., *Results and Problems in Cell Differentiation: MAP Kinases in Plant Signal Transduction* (ed. Hirt, H.) 85-93 (Springer, Heidelberg, 2000); Ligterink et al., *Science* 276:2054-2057, 1997; Eulgem et al., *EMBO J.* 18:4689-4699, 1999), the *Arabidopsis* mesophyll protoplast transient expression system offers a new tool to study defense signaling based on early gene transcription.

Example 9: Flg22 signals through FLS2 in *Arabidopsis* protoplasts

To determine whether the *WRKY29*, *FRK1*, and *GST1* promoters were activated by flg22 through the FLS2 LRR receptor kinase (Gomez-Gomez et al., *Mol. Cell* 5:1003-1011, 2000), protoplasts isolated from *fls2* mutant leaves were transiently transfected with the *WRKY29-LUC*, *FRK1-LUC*, and *GST1-LUC*,

reporter constructs. In the *fls2* mutant protoplasts, none of the three promoters could be activated by flg22 (Fig. 17C). Transient expression of wild-type FLS2 (Gomez-Gomez et al., *Mol. Cell* 5:1003-1011, 2000) restored flg22-inducibility of the promoters in the *fls2* mutant protoplasts (Fig. 17C). Consistent with these results, the broad-spectrum protein kinase inhibitor staurosporine (ST) effectively blocked the ability of flg22 to activate the *WRKY29*, *FRK1*, and *GST1* promoters (Fig. 17B). These data showed that flagellin signaling leading to the expression of *WRKY29*, *FRK1*, and *GST1* requires FLS2.

Example 10: Specific MAPKs in flg22 signaling

The role of MAPK signaling in the flg22-mediated activation of the *WRKY29*, *FRK1*, and *GST1* promoters was tested by using the MAPKK inhibitor U0126. As shown in Fig. 17B, U0126 partially blocked transcription of the *WRKY29* and *FRK1* promoters but had no effect on the *GST1* promoter. To further test the involvement of MAPK signaling in the activation of *WRKY29* and *FRK1* promoters, a mouse MAPK phosphatase (MKP1) was transiently expressed in *Arabidopsis* protoplasts (Kovtun et al., *Nature* 395:716-20, 1998). Similar to U0126, MKP1 partly reduced flg22-induced activation of *WRKY29* and *FRK1* promoters but had no effect on the *GST1* promoter (Fig. 17D). These results suggested that both MAPK-dependent and MAPK-independent signaling pathways act downstream of FLS2 to activate the *WRKY29* and *FRK1* promoters.

Since flg22 activation of *WRKY29* and *FRK1* promoters was partially blocked by U0126 and MKP1, we examined whether flg22 can activate MAPKs in *Arabidopsis* protoplasts. Treatment of protoplasts with flg22 resulted in rapid activation of endogenous protein kinases (PKs) that phosphorylated myelin basic protein (MBP), a commonly used MAPK substrate (Fig. 18A). These PKs were not activated by treatment with flg23RM (data not shown), suggesting their importance in pathogen defense. To systematically test the *Arabidopsis* MAPKs (MPKs) to determine which one(s) are involved in flg22 signaling, we chose six representative MPKs corresponding to four out of the five MPK subfamilies that may exhibit distinct functions based on sequence homology analysis (Tena et al., *Curr. Opin. Plant Biol.* 4:392-400, 2001; Mizoguchi et al, *Results and Problems in Cell Differentiation: MAP Kinases in Plant Signal Transduction* (ed. Hirt, H.) 29-38 (Springer, Heidelberg, 2000)). These MPKs were tagged with the haemagglutinin

(HA) epitope, transiently expressed in protoplasts, immunoprecipitated with an anti-HA antibody, and tested *in vitro* for kinase activity (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). The results indicated that MPK3 and MPK6, which belong to the same subfamily, but not the other MPKs, showed strong MBP phosphorylation following flg22 treatment (Fig. 18B). The involvement of MPK5 cannot be ruled out, however, due to its poor expression and/or instability in mesophyll protoplasts. The result is consistent with the previously reported activation of MPK6 in *Arabidopsis* cultured cells and leaf strips by flg22 (Nuhse et al., *J. Biol. Chem.* 275:7521-7526, 2000). The activation of MPK3 and MPK6 by flg22 did not occur in the *fls2* mutant protoplasts unless functional FLS2 was co-expressed (Fig. 18C). Importantly, when a kinase inactive mutant of FLS2 (FLS2Km) was expressed in the *fls2* mutant protoplasts, flg22-triggered activation of MPK3 and MPK6 was not observed (Fig. 18C), suggesting that the kinase activity of FLS2 is important for flagellin signaling and MAPK activation. Furthermore, the co-expression of MKP1, which partially blocked flg22-induced activation of *WRKY29* and *FRK1* promoters (Fig. 17D), eliminated MPK3 and MPK6 activation by flg22 (Fig. 18D). These results suggest that flagellin perception by FLS2 leads to the activation of MPK3 and/or MPK6. The studies also revealed the importance of MAPK-independent pathway in flg22 signaling.

20 Example 11: Redundant MAPKKs in flg22 signaling

The *Arabidopsis* genome contains nine MAPKK (MKK) genes that belong to four subfamilies, suggesting at least four distinct functions (Tena et al., *Curr. Opin. Plant Biol.* 4:392-400, 2001). Four of these MKKs, MKK1, MKK2, MKK4, and MKK5, belonging to two subfamilies, are expressed in leaf cells (Tena et al., *Curr. Opin. Plant Biol.* 4:392-400, 2001; Mizoguchi et al, *Results and Problems in Cell Differentiation: MAP Kinases in Plant Signal Transduction* (ed. Hirt, H.) pp. 29-38 (Springer, Heidelberg, 2000)). Therefore, these four MKK genes were cloned and analyzed in the protoplast transient expression assay. To distinguish MKKs from the HA-tagged MPKs in transfected protoplasts, a MYC-epitope tag was fused to each MKK. Since MAPKKs require phosphorylation to be activated, gain-of-function mutants of the four MKKs were generated by converting the conserved serine and/or threonine residues in the kinase-activation loop located between subdomains VII and VIII to aspartate or glutamate to mimic phosphorylation. The

MYC-tagged wild-type or constitutively active MKKs were co-expressed individually with the six HA-tagged MPKs to systematically determine their regulatory relationships. The results showed that constitutively active MKK4 and MKK5 (MKK4a and MKK5a, respectively), which belong to the same subfamily, were equally effective at phosphorylating (Fig. 20A) and activating MPK3 and MPK6 (Figs. 19A and 19B). We cannot rule out the possibility that MPK5 is also activated by MKK4a and/or MKK5a since the expression of MPK5 was poor (Fig. 19A). In contrast, constitutively active MKK1 and MKK2 (MKK1a and MKK2a, respectively) were unable to activate these MPKs (Fig. 19B), although MKK1a did activate other MPKs (data not shown).

To directly test the role of MKK4 and MKK5 in flg22 signaling, plasmid DNA expressing the constitutively active MKK constructs was cotransfected with various reporter gene constructs into protoplasts (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). Either MKK4a or MKK5a, but not MKK1a or MKK2a, was sufficient to specifically activate the *WRKY29* and *FRK1* promoters (Fig. 19C). To further test the importance of MKK4 and MKK5 in the early defense response, corresponding dominant-negative mutants were generated by replacing the conserved lysine residue in the ATP binding site with methionine. As shown in Fig. 19D, dominant-negative MKK4 or MKK5 but not MKK1 (data not shown) could partially block flg22 activation of the *WRKY29* and *FRK1* promoters similarly to U0126 or MKP1 (Figs. 17B and 17D). In addition, MKK4a and MKK5a, but not MKK1a and MKK2a, could bypass the requirement of FLS2 in flg22 signaling, demonstrated by their ability to activate the *WRKY29* and *FRK1* promoters in *fls2* mutant protoplasts (Fig. 19E).

Thus, on the basis of both gain-of-function and loss-of-function analyses, it appears that flagellin signaling activates MKK4 and MKK5, which in turn phosphorylate and activate MPK3 and MPK6, leading to the expression of early defense response genes. The functions of MKK4 and MKK5 in flg22 signaling are likely redundant in *Arabidopsis* leaf cells (see below). It has been proposed that MPK3 and/or MPK6 mediate expression of *GST6* in *Arabidopsis* protoplasts treated with H₂O₂ (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). A MAPKK functioning in this oxidative-stress signaling pathway has not been identified. Although MKK4 and MKK5 can activate MPK3 and MPK6, neither

MKK4a nor MKK5a was able to induce the expression of *GST6*, a reporter gene in the oxidative stress signaling pathway (Figs. 19C and 19E). This result suggests that flagellin and H₂O₂ may activate different MAPK signaling cascades.

Example 12: A specific MAPKKK in flg22 signaling

5 On the basis of sequence homology in the kinase domains to mammalian MAPKKKs, there are at least 25 putative *Arabidopsis* MAPKKKs that can be divided into six subgroups (Tena et al., *Curr. Opin. Plant Biol.* 4:392-400, 2001). The sequence similarity among the subgroups is low, especially in the regulatory domains, indicating that the subgroups have distinct functions (Tena et al., *Curr.*
 10 *Opin. Plant Biol.* 4:392-400, 2001; Jouannic et al., *Gene* 233:1-11, 1999). We analyzed four *Arabidopsis* MAPKKKs, CTR1, EDR1, MEKK1, and ANP1, from three subgroups expressed in leaf cells, to determine whether any of them could activate MKK5. Constitutively active derivatives of these four MAPKKKs were constructed by deleting the putative regulatory domain (Kovtun et al., *Proc. Natl.*
 15 *Acad. Sci., U.S.A.* 97:2940-2945, 2000). The four constitutively active MAPKKKs were fused to a HA-epitope tag and coexpressed individually with MYC-epitope-tagged wild-type MKK5 in protoplasts. To determine whether MKK5 had been phosphorylated and activated, MKK5 was immunoprecipitated using an anti-MYC antibody, and its activity was determined *in vitro* using a purified kinase inactive
 20 MPK6 fused to glutathione S-transferase (GST-MPK6Km) as a substrate. As shown in Fig. 20A, MKK5 was specifically activated by constitutively active MEKK1 (Δ MEKK1) and could phosphorylate MPK6. Constitutively active ANP1 (Δ ANP1), which can mimic the H₂O₂ signal and induce the *GST6* promoter (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000), activated MKK5 only
 25 marginally despite a high level of protein expression (Fig. 20A), supporting the idea that flagellin and H₂O₂ activate different MAPK pathways.

Consistent with the above results, Δ MEKK1 activated the *WRKY29* and *FRK1* promoters in the absence of flg22 in both wild-type and *fls2* mutant protoplasts (Figs. 20B and 20C) but not the activity of the H₂O₂- or Δ ANP1-
 30 inducible *GST6* promoter (Fig. 20B). Furthermore, constitutively active CTR1 (Δ CTR1), which did not activate MKK5, had no effect on the *WRKY29* and *FRK1* promoter activities (Figs. 20B and 20C). To further test the involvement of MEKK1 in flg22 signaling, a dominant-negative mutant was generated by altering the ATP

binding site but maintaining the regulatory and protein-protein interaction domains of the full-length MEKK1. The dominant-negative mutant (MEKK1ⁱⁿ) partially suppressed flg22 activation of *WRKY29* and *FRK1* promoters (Fig. 20D), similarly to U0126 treatment, MKP1, and dominant-negative MKK4 or MKK5 (Figs. 17B, 17D, and 19D). Dominant negative CTR1 did not have an effect (data not shown). Taken together, the results described so far strongly suggest that flagellin perception by FLS2 leads to the induction of *WRKY29* and *FRK1* transcription through the activation of a MAPK signaling cascade consisting of MEKK1, MKK4/MKK5, and MPK3/MPK6.

10 Example 13: Positive feedback control by WRKY transcription factors

WRKY proteins have been shown to bind to W box DNA elements (containing a TGAC core sequence) that are found in the promoters of many defense-related genes including *WRKY29* (10 W boxes) and *FRK1* (15 W boxes) (Maleck et al., *Nature Genet.* 26:403-410, 2000; Euglem et al., *Trends Plant Sci.* 5:199-206, 2000; Du et al., *Plant J.* 24:837-847, 2000). In parsley protoplasts, a fungal pathogen-inducible WRKY1 protein, with two WRKY domains, is targeted to the nucleus and activates its own promoter by binding to multiple W boxes (Eulgem et al., *EMBO J.* 18:4689-4699, 1999). *Arabidopsis* WRKY29, with a single WRKY domain, is not the ortholog of parsley WRKY1 (Eulgem et al., *EMBO J.* 18:4689-4699, 1999), and is the *WRKY* gene shown to be induced by pathogen-derived flagellin (Fig. 17). WRKY29-GFP, a WRKY29 and green fluorescent protein fusion, was constitutively localized in the nucleus in the presence or absence of flg22 (Fig. 21A). Transient expression of WRKY29 activated its own promoter in the absence of flg22 (Fig. 21B), indicating a positive feedback control (Eulgem et al., *EMBO J.* 18:4689-4699, 1999). WRKY29 also strongly activated the *FRK1* promoter, but suppressed the basal activities of the *GST6* and *RD29A* promoters (Fig. 21C). Transient expression of WRKY29 activated the *WRKY29* and *FRK1*, but not *RD29A*, promoters in the *fls2* mutant protoplasts, suggesting that WRKY29 acts downstream of the flagellin receptor (Fig. 21C). Treatment of WRKY29-expressing protoplasts with flg22 did not further enhance the activity of the *WRKY29* and *FRK1* promoters (Fig. 21B). These results are consistent with the idea that unmodified wild-type WRKY29 can act as a transcriptional activator. In contrast to WRKY29, a different *Arabidopsis* WRKY protein, WRKY42 (At4g04450), did not activate the

WRKY29 and FRK1 promoters, but rather slightly inhibited the flg22 activation of these promoters (Fig. 21B). Moreover, unlike WRKY29, WRKY42 did not suppress basal activities of the GST6 and RD29A promoters (Fig. 21B). Thus, we postulated that WRKY29 is a key transcriptional activator involved in the expression of defense genes in *Arabidopsis* innate immune responses. Similarly to MKK and MPK gene families, the large WRKY family also have members that are highly homologous to WRKY29. The homologous WRKY22 (At4g01250) in the same WRKY subgroup (Euglem et al., *Trends Plant Sci.* 5:199-206, 2000) was cloned and tested in the protoplast transient assay. WRKY22 regulated these promoters similarly to WRKY29 (Fig. 21B), suggesting that WRKY29 and WRKY22 may be functionally redundant.

Example 14: Resistance to bacterial and fungal pathogens

On the basis of the results obtained with the protoplast transient expression assays, we tested whether *Arabidopsis* plants exhibited enhanced resistance to bacterial pathogens when the flagellin MAPK cascade is constitutively activated or WRKY29 is expressed by means of *Agrobacterium*-mediated transient transformation (Xiang et al., *Plant Mol. Biol.* 40:711-717, 1999). This method has allowed a reliable assessment of the effect of transgenes in leaves (Nimchuk et al., *Cell* 101:353-363, 2000; Swiderski et al., *Plant J* 26:101-12, 2001; Yang et al., *Proc. Natl. Acad. Sci., U.S.A.* 98:741-746, 2001). When infected by the virulent bacterial pathogen *Pseudomonas syringae*, transiently transformed control leaves expressing GFP developed chlorotic lesions (Fig. 21D). Similar results were obtained with untransformed leaves and leaves expressing Δ CTR1, MKK1a, or wild-type WRKY42 as controls (not shown). In contrast, transformed leaves expressing Δ MEKK1, MKK4a, MKK5a or wild-type WRKY29 displayed enhanced resistance to *P. syringae* (Fig. 21D). The observation that both MKK4a and MKK5a can confer pathogen resistance is consistent with the idea that these MAPKKs are functionally redundant.

Importantly, development of soft-rot symptoms caused by infection with the fungal pathogen *Botrytis cinerea* was also effectively suppressed when Δ MEKK1, MKK4a, or WRKY29, but not GFP, was transiently expressed in leaves (Fig. 21E). Microscopic analysis indicated that in *Arabidopsis* leaves expressing MKK1a as a control, fungal spores germinated and formed superficial mycelium and branched

appressoria filled with intensely stained cytoplasm within two days after *B. cinerea* infection (Fig. 21F). In contrast, only germ tubes were formed during the same period of time in leaves expressing MKK4a. Similar results were obtained with Δ MEKK1, MKK5a, and WRKY29 (data not shown). These results suggest that

5 defense responses activated by the flagellin MAPK cascade or WRKY29 are specifically effective against fungal as well as bacterial pathogens.

In addition, transient expression of MKK4a in *Arabidopsis* leaves, as in Fig. 21D, inhibited the growth of *P. syringae maculicola* ES4326 15-fold one day after infiltration, and over 100-fold four days after infiltration. The results are consistent

10 with the leaf phenotype shown in Fig. 21D.

Examples 8-14 described above were performed using the following methods.

Reporter constructs

To identify early response genes, a subtracted cDNA library that represented

15 mRNA species induced by the elicitor flg22 in *Arabidopsis* mesophyll protoplasts was generated as previously described (Asai et al., *Plant Cell* 12:1823-1835, 2000). The DNA regions immediately upstream from the translation start sites of the *WRKY29*, *FRK1*, and *GST1* genes were amplified by PCR from *Arabidopsis* (Col-0) genomic DNA. The sizes of the amplified fragments were 2.6 kb, 2.8 kb, and 0.9

20 kb, respectively. These promoters were fused to a luciferase (LUC) reporter gene to generate *WRKY29-LUC*, *FRK1-LUC*, and *GST1-LUC* as described by Kovtun et al., (*Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000).

Effector constructs

Arabidopsis cDNAs, encoding MAPKs: MPK2 (At1g59580), MPK3

25 (At3g45640), MPK5 (At4g11330), MPK6 (At2g43790), MPK7 (At2g18170), MPK9 (At3g18040); MAPKKs: MEK1 (At4g26070), MKK2 (At4g29810), MKK4 (At1g51660), MKK5 (At3g21220); MAPKKKs: CTR1 (At5g03730), EDR1 (At1g08720), MEKK1 (At4g08500), ANP1 (At1g09000); receptor FLS2 (At5g46330); transcription factors WRKY22 (At4g01250), WRKY29 (At4g23550)

30 and WRKY42 (At4g04450), were obtained by PCR from an *Arabidopsis* cDNA library and verified by sequencing. Mouse MKP1 phosphatase was a gift from H. Sun and N. Tonks (Kovtun et al., *Nature* 395:716-20, 1998). PCR products were

fused to the double HA (MAPK, MAPKKK, receptor and WRKY) or MYC (MAPKK and MKP1) epitope tag sequence and inserted into a plant expression vector or pCB302 containing the *35SC4PPDK* promoter and the *NOS* terminator (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000; Kovtun et al.,
 5 *Nature* 395:716-20, 1998; and Xiang et al., *Plant Mol. Biol.* 40:711-717, 1999).

The constitutively active forms of MAPKKs were generated by site-specific mutation, replacing the serine or threonine residues in the activation loop domain [S/T]XXXXX[S/T] by acidic amino acids glutamate or aspartate: MKK1(T218ES224D), MKK2(T220DT226E), MKK4(T224DS230E) and
 10 MKK5(T215ES221E). The active forms of MAPKKK were generated by keeping the catalytic domain only: Δ CTR1(544-799), Δ EDR1(662-921), Δ MEKK1(326-592), Δ ANP1(57-338). The inactive forms of all kinases were generated by site-specific mutation, replacing the conserved lysine residues in the kinase ATP binding loop by a methionine: FLS2(K898M), MKK4(K108M), MKK5(K99M),
 15 MPK6(K92M), MEKK1(K361M). All these mutations were verified by sequencing.

GST fusion proteins

The GST fusions were generated by subcloning the coding sequences of MAPKs and MAPKKs into pGEX-4T-1 vector (Amersham Pharmacia Biotech) in frame with the GST coding sequence. All the GST fusion proteins were purified
 20 based on the manufacture's procedure and subjected to *in vitro* phosphorylation assay. Except the kinase inactive mutants that were used as controls, all PKs showed significant autophosphorylation activity and were able to phosphorylate an exogenous substrate, indicating that they are active kinases (data not shown).

Arabidopsis mesophyll protoplast transient expression assay

25 Protoplast transient expression assay was carried out as described previously (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). Promoter activities are represented by LUC/GUS activities, and normalized to the value obtained with protoplasts without the treatments. In case of protoplasts prepared from *fls2* mutant plants, promoter activities are normalized to the value obtained
 30 with wild-type protoplasts transfected with a control GFP plasmid. Constitutively active MAPKKKs, immunoprecipitated from transfected protoplasts, phosphorylated casein *in vitro* (Kovtun et al., *Nature* 395:716-20, 1998), indicating that they are active kinases. GFP fluorescence was observed by either Nikon TE200

fluorescent microscopy or with Leica TCSNT Confocal Spectrophotometer (Germany). Coexpression of a nuclear-targeted red-fluorescent protein was used as a control (Patharkar et al., *Plant J.* 24:679-691, 2000).

RT-PCR

5 Samples were taken at indicated time points after treatment and total RNA was prepared as described (Asai et al., *Plant Cell* 12:1823-1835, 2000). RT-PCR was performed with 1 ng of total RNA and 0.6 μ M of each primer using OneStep RT-PCR Kit (Qiagen, Germany). PCR was run for 35 cycles. Constitutively expressed *UBQ10* (AT4g05320) or *ACTIN-1* (At2g37620) mRNA was always co-
10 amplified with each mRNA and used as an internal standard. For these genes, 0.1 μ M of each primer was used. Typical results for *UBQ10* mRNA are shown.

Protein Kinase assays

The MAPK in-gel kinase assay was carried out as described by Kovtun et al. (*Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). For immuno-complex assays,
15 tagged kinases were immunoprecipitated from lysates of transfected protoplasts with the corresponding antibody and analyzed with a known substrate as indicated (MBP or GST-MPK6 Km).

Agrobacterium-mediated transient transformation

To analyze plant susceptibility to a bacterial pathogen, five-week-old
20 *Arabidopsis* plants were infiltrated with a suspension (containing 100 μ M acetosyringone) of *A. tumefaciens* carrying a binary vector plasmid pCB302 (Xiang et al., *Plant Mol. Biol.* 40:711-717, 1999) expressing GFP, Δ MEKK1, MKK4a, MKK5a, or WRKY29. The plants were incubated for three days under the conditions as described (Asai et al., *Plant Cell* 12:1823-1835, 2000). Infiltrated
25 leaves were then inoculated with 10 mM MgSO_4 or the same solution with suspended *P. syringae maculicola* ES4326 (10^4 cfu/cm²) at the location of transient gene expression. The leaves were photographed three days later.

To analyze plant susceptibility to a fungal pathogen, a suspension (containing 100 μ M acetosyringone and 0.01% Silwett L-77) of *A. tumefaciens*
30 carrying a binary vector plasmid expressing GFP, Δ MEKK1, MKK4a, MKK1a, or WRKY29 was first applied to the lower surface of a leaf of six-week-old *Arabidopsis*. The leaves were incubated for 24 hours and then the upper surface of

transformed leaves was infected with *B. cinerea* sclerotia. The infected leaves were photographed five days later after removing the fungal sclerotia. To observe the early stage of *B. cinerea* development, 5 µl drops of a 5×10^5 spores/ml suspension were placed on the upper surface of an *Arabidopsis* leaf. Two days after infection, fungal structures were visualized by Trypan blue-lactophenol staining and observed with Confocal Spectrophotometer TCSNT (Leica, Germany). Untransformed and GFP-expressing control leaves developed similar lesions as the MKK1a-expressing leaves.

10 Summary

We have demonstrated that *Arabidopsis* mesophyll protoplast transient expression assays can be combined with genetic and genomic information to provide a powerful tool for the analysis of MAPK signaling involved in plant innate immunity. Importantly, the key components of the innate immune signaling pathway identified using the leaf cell system confer pathogen resistance when expressed in leaves of intact plants (Fig. 21). Since wild-type WRKY22 or WRKY29 is sufficient to mimic flg22 and MAPK signaling, we postulate that a specific WRKY inhibitor may be phosphorylated and inactivated by the flagellin MAPK cascade upon pathogen infection. As summarized in the model shown in Figure 22, this would make the flg22 signaling pathway partially analogous to signaling pathways in insects and mammals, in which TLRs initiate a cascade of signaling events leading to phosphorylation and degradation of an inhibitor of a Rel-like transcription factor (Aderem et al., *Nature* 406:782-787, 2000; Khush et al., *Trends Genet.* 16:442-449, 2000). It will be interesting to test whether the flg22- and FLS2-dependent phosphorylation of the ankyrin-containing protein AtPhos43, recently identified by a proteomic approach (Peck et al, *Plant Cell* 13:1467-1475, 2001), is MAPK dependent and can control WRKY22 and WRKY29 sub-cellular localization and/or activity.

We further propose that early signaling events initiated by diverse pathogens converge into a conserved MAPK cascade that positively regulates plant defense responses (Fig. 22). This is consistent with the observation that similar plant defense responses are elicited by many viral, bacterial, and fungal pathogens. Because MAPK signaling components are highly conserved among diverse plant

species, our results with flagellin elicitation provides insights into the genetic manipulation of resistance to a wide array of pathogens in economically important crop plants.

As shown in Fig. 17B, 17D, 19D and 20D, our data also reveal the previously unexpected existence of a MAPK-independent pathway in flg22 signaling.

The lack of MAPK cascade mutations in the flg22 signaling pathway is most likely due to functional redundancy in the pathway, as suggested by the identification of the MKK4/MKK5 and MPK3/MPK6 pairs in our study. The WRKY22 and WRKY29 transcription factors in the flg22 signaling pathway may also provide redundant functions. Furthermore, similar to the yeast MAPKKK STE11 (Madhani et al., *Trends Genet.* 14:151-155, 1998), some plant MAPK signaling components could be involved in more than one pathway (Tena et al., *Curr. Opin. Plant Biol.* 4:392-400, 2001). The H₂O₂-activated MAPK cascade shares MPK3 and MPK6 with the flg22 pathway but activates different target genes (Kovtun et al., *Proc. Natl. Acad. Sci., U.S.A.* 97:2940-2945, 2000). The MAPKKK of the flagellin cascade, MEKK1, may also initiate a different MAPK cascade. In addition, it should be noted that, besides the flagellin MAPK cascade, other pathways utilizing untested MAPK, MAPKK, and MAPKKK genes could also play important roles in *Arabidopsis* innate immunity. Finally, plants transiently expressing ΔMEKK1, MKK4a, or WRKY29 are resistant to both bacterial and fungal pathogens, suggesting that other pathogen-derived signals in addition to flg22 may activate the MEKK1, MKK4/MKK5, MPK3/MPK6, WRKY22/29 signaling pathway through distinct receptors (Fig. 22) (Peck et al, *Plant Cell* 13:1467-1475, 2001). Given this complexity, a combination of genetic, genomic, cellular, and biochemical approaches including the construction of knock-out mutants and expression of signaling components in transgenic plants will be required to untangle the intertwined signaling webs mediated by MAPK cascades in plants.

Methods for isolating sequences encoding MAPKKKs, MAPKKs, and WRKYs now follow. In addition, methods for expressing such sequences in a plant and methods for engineering disease resistant plants are also described.

Isolation of Sequences Encoding MAPKKKs, MAPKKs, and WRKYs

The isolation of additional MAPKKK (MAPKKs and WRKYs) coding sequences, as well as MAPKKK and MAPKK kinase domains, having the ability to regulate pathogen defense in plants is accomplished using standard strategies and techniques that are well known in the art.

In one particular example, the tobacco NPK1 sequences (or, for example, Arabidopsis ANP1, ANP2, or ANP3 sequences) may be used, together with conventional screening methods of nucleic acid hybridization screening, to isolate additional sequences encoding MAPKKK polypeptides (or kinase domain-containing fragments thereof), as well as kinase domains of MAPKKK. Such hybridization techniques and screening procedures are well known to those skilled in the art and are described, for example, in Benton and Davis, *Science* 196: 180, 1977; Grunstein and Hogness, *Proc. Natl. Acad. Sci., USA* 72: 3961, 1975; Ausubel et al. *Current Protocols in Molecular Biology*, Wiley Interscience, New York; Berger and Kimmel, *Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York.; and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York. In one particular example, all or part of the NPK1 gene may be used as a probe to screen a recombinant plant DNA library for genes having sequence identity or similarity to the NPK1 gene or its kinase domain. Hybridizing sequences are detected by plaque or colony hybridization according to the methods described below.

Alternatively, using all or a portion of the amino acid sequence of the kinase domain, one may readily design kinase domain-specific oligonucleotide probes, including kinase domain degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either DNA strand and any appropriate portion of the kinase domain sequence. General methods for designing and preparing such probes are provided, for example, in Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience, New York; and Berger and Kimmel, *Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York. These oligonucleotides are useful for kinase domain sequence isolation, either through their use as probes capable of hybridizing to kinase complementary sequences or as primers for various amplification techniques, for example,

polymerase chain reaction (PCR) cloning strategies. If desired, a combination of different oligonucleotide probes may be used for the screening of a recombinant DNA library. The oligonucleotides may be detectably-labeled using methods known in the art and used to probe filter replicas from a recombinant DNA library.

- 5 Recombinant DNA libraries are prepared according to methods well known in the art, for example, as described in Ausubel et al. (supra), or they may be obtained from commercial sources.

As discussed above, kinase domain-specific oligonucleotides may also be used as primers in amplification cloning strategies, for example, using PCR. PCR
10 methods are well known in the art and are described, for example, in *PCR Technology*, Erlich, ed., Stockton Press, London, 1989; *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, Inc., New York, 1990; and Ausubel et al. (supra). Primers are optionally designed to allow cloning of the amplified product into a suitable vector, for example, by including appropriate
15 restriction sites at the 5' and 3' ends of the amplified fragment (as described herein). If desired, kinase domain sequences may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al. (supra)). By this method, oligonucleotide primers based on an kinase domain sequence are oriented in the 3' and 5' directions and are used to generate overlapping PCR
20 fragments. These overlapping 3'- and 5'-end RACE products are combined to produce an intact full-length cDNA. This method is described in Innis et al. (supra); and Frohman et al., *Proc. Natl. Acad. Sci., USA* 85: 8998, 1988.

Confirmation of a sequence's relatedness to the kinase domains of the NPK and ANP MAPKKKs may be accomplished by a variety of conventional methods
25 including, but not limited to, sequence comparison of the gene and its expressed product. In addition, the activity of the gene product may be evaluated according to any of the techniques described.

Once a MAPKKK gene or its kinase domain (or a gene encoding a MAPKK or WRKY polypeptide) is identified, it is cloned according to standard methods and
30 used for the construction of plant expression vectors as described below.

Expression Constructs

Those skilled in the field of molecular biology will understand that any of a wide variety of expression systems may be used to provide the recombinant protein. The precise host cell used is not critical to the invention. A MAPKKK polypeptide or its kinase domain (or a MAPKK or WRKY polypeptide) may be produced in a prokaryotic host, for example, *E. coli*, or in a eukaryotic host, for example, *Saccharomyces cerevisiae*, mammalian cells (for example, COS 1 or NIH 3T3 cells), or any of a number of plant hosts including, without limitation, algae, tree species, ornamental species, temperate fruit species, tropical fruit species, vegetable species, legume species, crucifer species, monocots, dicots, or in any plant of commercial or agricultural significance. Particular examples of suitable plant hosts include, but are not limited to, Conifers, Petunia, Tomato, Potato, Tobacco, Arabidopsis, Lettuce, Sunflower, Oilseed rape, Flax, Cotton, Sugarbeet, Celery, Soybean, Alfalfa, Medicago, Lotus, Vigna, Cucumber, Carrot, Eggplant, Cauliflower, Horseradish, Morning Glory, Poplar, Walnut, Apple, Grape, Asparagus, Rice, Maize, Millet, Onion, Barley, Orchard grass, Oat, Rye, and Wheat. In addition, as is discussed below, expression constructs may be expressed in a transgenic plant to turn on the pathogen defense MAPK signal transduction pathway to enhance plant tolerance to its pathogen(s).

Materials for expressing these genes are available from a wide range of sources including the American Type Culture Collection (Rockland, MD); or from any of a number seed companies, for example, W. Atlee Burpee Seed Co. (Warminster, PA), Park Seed Co. (Greenwood, SC), Johnny Seed Co. (Albion, ME), or Northrup King Seeds (Harstville, SC). Descriptions and sources of useful host cells are also found in Vasil I.K., *Cell Culture and Somatic Cell Genetics of Plants*, Vol I, II, III *Laboratory Procedures and Their Applications*, Academic Press, New York, 1984; Dixon, R.A., *Plant Cell Culture-A Practical Approach*, IRL Press, Oxford University, 1985; Green et al., *Plant Tissue and Cell Culture*, Academic Press, New York, 1987; and Gasser and Fraley, *Science* 244: 1293, 1989.

The method of transformation or transfection and the choice of vehicle for expression of the MAPKKK polypeptide or its kinase domain (or a MAPKK or WRKY polypeptide) will depend on the host system selected. Transformation and transfection methods are described, e.g., in Ausubel et al. (supra); Weissbach and

Weissbach, *Methods for Plant Molecular Biology*, Academic Press, 1989; Gelvin et al., *Plant Molecular Biology Manual*, Kluwer Academic Publishers, 1990; Kindle, K., Proc. Natl. Acad. Sci., U.S.A 87: 1228, 1990; Potrykus, I., *Annu. Rev. Plant Physiol. Plant Mol. Biology* 42: 205, 1991; and BioRad (Hercules, CA) Technical Bulletin #1687 (Biolistic Particle Delivery Systems). Expression vehicles may be chosen from those provided, e.g., in *Cloning Vectors: A Laboratory Manual* (P.H. Pouwels et al., 1985, Supp. 1987); Gasser and Fraley (supra); Clontech Molecular Biology Catalog (Catalog 1992/93 Tools for the Molecular Biologist, Palo Alto, CA); and the references cited above. Other expression constructs are described by Fraley et al. (U.S. Pat. No. 5,352,605).

Most preferably, a MAPKKK polypeptide or its kinase domain (or a MAPKK or WRKY polypeptide) is produced by a stably-transfected plant cell line, a transiently-transfected plant cell line, or by a transgenic plant. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in Pouwels et al. (supra), Weissbach and Weissbach (supra), and Gelvin et al. (supra). Methods for constructing such cell lines are described in, e.g., Weissbach and Weissbach (supra), and Gelvin et al. (supra). Typically, plant expression vectors include (1) a cloned plant gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (for example, one conferring inducible or constitutive, pathogen- or wound-induced, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

Once the desired nucleic acid sequence encoding a MAPKKK polypeptide or its kinase domain (or a MAPKK or WRKY polypeptide) is obtained as described above, it may be manipulated in a variety of ways known in the art. For example, where the sequence involves non-coding flanking regions, the flanking regions may be subjected to mutagenesis.

For example, the kinase domain sequence (or a MAPKKK polypeptide or kinase domain-containing fragment thereof), if desired, may be combined with other DNA sequences in a variety of ways. Such a sequence may be employed with all or

part of the gene sequences normally associated with itself. In its component parts, a DNA sequence encoding a MAPKKK polypeptide or its kinase domain is combined in a DNA construct having a transcription initiation control region capable of promoting transcription and translation in a host cell.

5 In general, the constructs will involve regulatory regions functional in plants which provide for modified production of the regulator protein as discussed herein. The open reading frame coding for the regulator protein or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region such as the sequence naturally found in the 5' upstream region of the MAPKKK
10 polypeptide or its kinase domain; a MAPKK; or a WRKY polypeptide. Numerous other transcription initiation regions are available which provide for constitutive or inducible regulation.

For applications where developmental, cell, tissue, hormonal, or environmental expression is desired, appropriate 5' upstream non-coding regions are
15 obtained from other genes, for example, from genes regulated during meristem development, seed development, embryo development, or leaf development.

Regulatory transcript termination regions may also be provided in DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the MAPKKK polypeptide (or a MAPKK or
20 WRKY polypeptide) or any convenient transcription termination region derived from a different gene source. The transcript termination region will contain preferably at least 1-3 kb of sequence 3' to the structural gene from which the termination region is derived. Plant expression constructs having, for example, a MAPKKK protein kinase domain (e.g., the NPK1 kinase domain); a MAPKK
25 polypeptide; or a WRKY polypeptide as the DNA sequence of interest for expression may be employed with a wide variety of plant life. Such genetically-engineered plants are useful for a variety of industrial and agricultural applications as discussed herein. Importantly, this invention is applicable to dicotyledons and monocotyledons, and will be readily applicable to any new or improved
30 transformation or regeneration method.

An example of a useful plant promoter according to the invention is a caulimovirus promoter, for example, a cauliflower mosaic virus (CaMV) promoter. These promoters confer high levels of expression in most plant tissues, and the

activity of these promoters is not dependent on virally encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odell et al., *Nature* 313: 810, 1985). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., *Plant Cell* 2: 591, 1990; Terada and Shimamoto, *Mol. Gen. Genet.* 220: 389, 1990). Moreover, activity of this promoter can be further increased (i.e., between 2-10 fold) by duplication of the CaMV 35S promoter (see e.g., Kay et al., *Science* 236: 1299, 1987; Ow et al., *Proc. Natl. Acad. Sci., U.S.A.* 84: 4870, 1987; and Fang et al., *Plant Cell* 1: 141, 1989). In addition, the a minimal 35S promoter may also be used as is described herein.

Other useful plant promoters include, without limitation, the nopaline synthase promoter (An et al., *Plant Physiol.* 88: 547, 1988) and the octopine synthase promoter (Fromm et al., *Plant Cell* 1: 977, 1989).

For certain applications, it may be desirable to produce the MAPKKK polypeptide or its kinase domain (or a MAPKK or WRKY polypeptide) in an appropriate tissue, at an appropriate level, or at an appropriate developmental time. For this purpose, there are an assortment of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, shown to be regulated in response to the environment, hormones, and/or developmental cues. These include gene promoters that are responsible for heat-regulated gene expression (see, e.g., Callis et al., *Plant Physiol.* 88: 965, 1988; Takahashi and Komeda, *Mol. Gen. Genet.* 219: 365, 1989; and Takahashi et al., *Plant J.* 2: 751, 1992), light-regulated gene expression (e.g., the Arabidopsis Cab2 photosynthetic, leaf specific promoter described by Mitra et al., *Plant Mol. Biol.* 12: 169-179, 1989; the pea rbcS-3A described by Kuhlemeier et al., *Plant Cell* 1: 471, 1989; the maize rbcS promoter described by Schöffner and Sheen, *Plant Cell* 3: 997, 1991; or the chlorophyll a/b-binding protein gene found in pea described by Simpson et al., *EMBO J.* 4: 2723, 1985), hormone-regulated gene expression (for example, the abscisic acid (ABA) responsive sequences from the Em gene of wheat described by Marcotte et al., *Plant Cell* 1: 969, 1989; the ABA-inducible HVA1 and HVA22, and rd29A promoters described for barley and *Arabidopsis* by Straub et al., *Plant Cell* 6: 617, 1994, Shen et al., *Plant Cell* 7: 295, 1995; and wound-induced gene expression (for example, of wunI described by Siebertz et al., *Plant Cell* 1: 961, 1989), organ-specific gene

expression (for example, of the tuber-specific storage protein gene described by Roshal et al., *EMBO J.* 6: 1155, 1987; the 23-kDa zein gene from maize described by Schernthaner et al., *EMBO J.* 7: 1249, 1988; or the French bean β -phaseolin gene described by Bustos et al., *Plant Cell* 1: 839, 1989; the vegetative storage protein promoter (soybean vspB) described by Sadka et al (*Plant Cell* 6: 737-749, 1994)), cycling promoters (e.g., the *Arabidopsis* cdc2a promoter described by Hemerly et al., *Proc. Natl. Acad. Sci., USA* 89: 3295-3299, 1992), senescence-specific promoters (e.g., the *Arabidopsis* SAG12 promoter described by Gan et al, *Science*: 270, 1986-1988, 1995), seed-specific promoters (for example, endosperm-specific or embryo-specific promoters), or pathogen-inducible promoters (for example, PR-1 or β -1,3 glucanase promoters).

Plant expression vectors may also optionally include RNA processing signals, e.g, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., *Genes and Dev.* 1: 1183, 1987). The location of the RNA splice sequences can dramatically influence the level of transgene expression in plants. In view of this fact, an intron may be positioned upstream or downstream of a MAPKKK polypeptide or its kinase-domain encoding sequence (or a MAPKK polypeptide encoding sequence or a WRKY polypeptide encoding sequence) in the transgene to modulate levels of gene expression.

In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., *Proc. Natl. Acad. Sci., U.S.A.* 84: 744, 1987; An et al., *Plant Cell* 1: 115, 1989). For example, the 3' terminator region may be included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly used terminators are derived from the octopine or nopaline synthase signals.

The plant expression vector also typically contains a dominant selectable marker gene used to identify those cells that have become transformed. Useful selectable genes for plant systems include genes encoding antibiotic resistance genes, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin, or spectinomycin. Genes required for photosynthesis may also be used as selectable markers in photosynthetic-deficient

strains. Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the bar gene encoding the enzyme phosphinothricin acetyltransferase and conferring resistance to the broad spectrum herbicide Basta® (Hoechst AG, Frankfurt, Germany).

- 5 Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the transformed cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/mL (kanamycin), 20-50 µg/mL
- 10 (hygromycin), or 5-10 µg/mL (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., by Vasil et al., supra.

It should be readily apparent to one skilled in the art of molecular biology, especially in the field of plant molecular biology, that the level of gene expression is dependent, not only on the combination of promoters, RNA processing signals, and terminator elements, but also on how these elements are used to increase the levels

15 of selectable marker gene expression.

Plant Transformation

- Upon construction of the plant expression vector, several standard methods are available for introduction of the vector into a plant host, thereby generating a
- 20 transgenic plant. These methods include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller, In: *Genetic Engineering*, vol 6, PWJ Rigby, ed, London, Academic Press, 1987; and Lichtenstein, C.P., and Draper, J., In: *DNA Cloning*, Vol II, D.M. Glover, ed, Oxford, IRI Press, 1985)), (2) the particle delivery system (see, e.g., Gordon-Kamm
- 25 et al., *Plant Cell* 2: 603,1990); or BioRad Technical Bulletin 1687, supra), (3) microinjection protocols (see, e.g., Green et al., supra), (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol.* 23: 451, 1982; or e.g., Zhang and Wu, *Theor. Appl. Genet.* 76: 835, 1988), (5) liposome-mediated DNA uptake (see, e.g., Freeman et al., *Plant Cell Physiol.* 25: 1353, 1984), (6)
- 30 electroporation protocols (see, e.g., Gelvin et al., supra; Dekeyser et al., supra; Fromm et al., *Nature* 319: 791, 1986; Sheen, *Plant Cell* 2: 1027, 1990; or Jang and Sheen, *Plant Cell* 6: 1665, 1994), and (7) the vortexing method (see, e.g., Kindle supra). The method of transformation is not critical to the invention. Any method

which provides for efficient transformation may be employed. As newer methods are available to transform crops or other host cells, they may be directly applied.

The following is an example outlining one particular technique, an *Agrobacterium*-mediated plant transformation. By this technique, the general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, cloning and DNA modification steps are carried out in *E. coli*, and the plasmid containing the gene construct of interest is transferred by conjugation or electroporation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria, for example, streptomycin, and another that will function in plants, for example, a gene encoding kanamycin resistance or herbicide resistance. Also present on the vector are restriction endonuclease sites for the addition of one or more transgenes and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the DNA region that will be transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad) used for the shooting, a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to pass through. As a result, the plastic macroprojectile smashes against the stopping plate, and the tungsten microprojectiles continue toward their target through the hole in the plate. For the instant invention the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

In general, transfer and expression of transgenes in plant cells are now routine practices to those skilled in the art, and have become major tools to carry out gene expression studies in plants and to produce improved plant varieties of agricultural or commercial interest.

5

Transgenic Plant Regeneration

Plant cells transformed with a plant expression vector can be regenerated, for example, from single cells, callus tissue, or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil *supra*; Green et al., *supra*; Weissbach and Weissbach, *supra*; and Gelvin et al., *supra*.

In one particular example, a cloned constitutively active kinase domain of a MAPKKK (or a MAPKKK polypeptide or a kinase-containing fragment thereof); a constitutively active MAPKK polypeptide; or a WRKY polypeptide; or any combination thereof, is placed under the control of the *nos* promoter and the nopaline synthase terminator and carrying a selectable marker (for example, kanamycin resistance) is transformed into *Agrobacterium*. Transformation of leaf discs (for example, of tobacco or potato leaf discs), with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (*Science* 227: 1229, 1985). Putative transformants are selected after a few weeks (for example, 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g. 100 µg/mL). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth. If desired, seeds from self-fertilized transgenic plants can then be sowed in a soil-less medium and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surfaced sterilized seeds on hormone-free kanamycin-containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, for example, Ausubel et al. *supra*; Gelvin et al. *supra*).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA by standard immunoblot and DNA detection techniques. Each positive transgenic plant and its transgenic progeny are unique in

comparison to other transgenic plants established with the same transgene.

Integration of the transgene DNA into the plant genomic DNA is in most cases random, and the site of integration can profoundly affect the levels and the tissue and developmental patterns of transgene expression. Consequently, a number of
5 transgenic lines are usually screened for each transgene to identify and select plants with the most appropriate expression profiles.

Transgenic lines are evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis are employed and include
10 PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al., supra). The RNA-positive plants are then analyzed for protein expression by Western immunoblot analysis using specific antibodies (see, e.g., Ausubel et al., supra). In addition, in situ hybridization and
15 immunocytochemistry according to standard protocols can be done using transgene-specific nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

In addition, if desired, once the recombinant MAPKKK polypeptide or its kinase domain, a MAPKK polypeptide, or a WRKY polypeptide, or any
20 combination thereof is expressed in any cell or in a transgenic plant (for example, as described above), it may be isolated, e.g., using affinity chromatography. In one example, an anti-MAPKKK polypeptide antibody (e.g., produced as described in Ausubel et al., supra, or by any standard technique) may be attached to a column and used to isolate the polypeptide. Lysis and fractionation of MAPKKK-producing
25 cells prior to affinity chromatography may be performed by standard methods (see, e.g., Ausubel et al., supra). Once isolated, the recombinant protein can, if desired, be further purified, for example, by high performance liquid chromatography (see, e.g., Fisher, Laboratory Techniques In Biochemistry And Molecular Biology, eds., Work and Burdon, Elsevier, 1980).

30

Engineering Disease Resistance

As discussed above, plasmid constructs designed for the expression of constitutively active MAPKKKs or MAPKKs or expression of a WRKY gene

product are useful, for example, for activating plant defense pathways that confer anti-pathogenic properties to a transgenic plant. Such genes are isolated from a host plant and may be engineered for expression in the same plant, a closely related species, or a distantly related plant species. For example, cruciferous genes may be engineered for constitutive low level expression and then transformed into an *Arabidopsis* host plant. Alternatively, the isolated cruciferous gene may be engineered for expression in other cruciferous plants, such as the Brassicas (for example, broccoli, cabbage, and cauliflower). Similarly, the MAPKKK, MAPKK, or WRKY is useful for expression in related solanaceous plants, such as tomato, potato, and pepper. To achieve pathogen resistance, it is important to express a constitutively active MAPKKK, MAPKK or a WRKY protein at an effective level. Evaluation of the level of pathogen protection conferred to a plant by ectopic expression of such polypeptides is determined according to conventional methods and assays.

In one working example, constitutive ectopic expression of the ANPI or MEKKK gene in Russet Burbank potato is used to control *Phytophthora infestans* infection. In one particular example, a plant expression vector is constructed that contains the appropriate cDNA sequence expressed under the control of the enhanced CaMV 35S promoter as described by McPherson and Kay (U.S. Patent 5,359,142). This expression vector is then used to transform Russet Burbank according to the methods described in Fischhoff et al. (U.S. Patent 5,500,365). To assess resistance to fungal infection, transformed Russet Burbank and appropriate controls are grown to approximately eight-weeks-old, and leaves (for example, the second or third from the top of the plant) are inoculated with a mycelial suspension of *P. infestans*. Plugs of *P. infestans* mycelia are inoculated on each side of the leaf midvein. Plants are subsequently incubated in a growth chamber at 27°C with constant fluorescent light.

Leaves of transformed Russet Burbank and control plants are then evaluated for resistance to *P. infestans* infection according to conventional experimental methods. For this evaluation, the number of lesions per leaf and percentage of leaf area infected are recorded every twenty-four hours for seven days after inoculation. From these data, levels of resistance to *P. infestans* are determined. Transformed

potato plants that express an constitutively active ANP1 or MEKK1 or both having an increased level of resistance to *P. infestans* relative to control plants are taken as being useful in the invention.

Alternatively, to assess resistance at the whole plant level, transformed and control plants are transplanted to potting soil containing an inoculum of *P. infestans*. Plants are then evaluated for symptoms of fungal infection (for example, wilting or decayed leaves) over a period of time lasting from several days to weeks. Again, transformed potato plants expressing the ANP1 or MEKK1 or both having an increased level of resistance to the fungal pathogen, *P. infestans*, relative to control plants are taken as being useful in the invention.

In another working example, expression of an MAPKKK, MAPKK, or WRKY in tomato is used to control bacterial infection, for example, to *Pseudomonas syringae*. Specifically, a plant expression vector is constructed that contains the cDNA sequence of one or more of the genes encoding these polypeptides is expressed under the control of the enhanced CaMV 35S promoter as described by McPherson and Kay, supra. This expression vector is then used to transform tomato plants according to the methods described in Fischhoff et al., supra. To assess resistance to bacterial infection, transformed tomato plants and appropriate controls are grown, and their leaves are inoculated with a suspension of *P. syringae* according to standard methods, for example, those described herein. Plants are subsequently incubated in a growth chamber, and the inoculated leaves are subsequently analyzed for signs of disease resistance according to standard methods. For example, the number of chlorotic lesions per leaf and percentage of leaf area infected are recorded and evaluated after inoculation. From a statistical analysis of these data, levels of resistance to *P. syringae* are determined. Transformed tomato plants that express an MAPKKK, MAPKK, or WRKY gene having an increased level of resistance to *P. syringae* relative to control plants are taken as being useful in the invention.

In still another working example, expression of an MAPKKK, MAPKK, or WRKY gene of rice is used to control fungal diseases, for example, the infection of tissue by *Magnaporthe grisea*, the cause of rice blast. In one particular approach, a plant expression vector is constructed that contains the cDNA sequence of the rice MAPKKK, MAPKK, or WRKY gene that is constitutively expressed under the

control of the rice actin promoter described by Wu et al. (WO 91/09948). This expression vector is then used to transform rice plants according to conventional methods, for example, using the methods described in Hiei et al. (*Plant Journal* 6:271-282, 1994). To assess resistance to fungal infection, transformed rice plants and appropriate controls are grown, and their leaves are inoculated with a mycelial suspension of *M. grisea* according to standard methods. Plants are subsequently incubated in a growth chamber and the inoculated leaves are subsequently analyzed for disease resistance according to standard methods. For example, the number of lesions per leaf and percentage of leaf area infected are recorded and evaluated after inoculation. From a statistical analysis of these data, levels of resistance to *M. grisea* are determined. Transformed rice plants that express a gene having an increased level of resistance to *M. grisea* relative to control plants are taken as being useful in the invention.

15 Use

The invention described herein is useful for a variety of agricultural and commercial purposes including, but not limited to, improving pathogen resistance, increasing crop yields, improving crop and ornamental quality, and reducing agricultural production costs. In particular, ectopic expression of a kinase domain of a MAPKKK polypeptide (or a MAPKKK polypeptide or a kinase domain-containing fragment thereof); a MAPKK polypeptide, or a WRKY polypeptide in a plant cell provides resistance to a plant pathogen and can be used to protect plants from pathogens that reduces plant productivity and viability. The invention therefore provides pathogen and disease resistance to to plants, especially crop plants, most especially crop plants such as tomato, potato, cotton, pepper, maize, wheat, rice, and legumes such as soybean, or any crop plant that is susceptible to pathogens. For example, transgenic maize and soybean may be genetically engineered to express a kinase domain of a MAPKKK (e.g., NPK1 or an ANP such as ANP1, ANP2, or ANP3), a MAPKK polypeptide, or a WRKY polypeptide, or any combination thereof according to standard methods, such as those described in Adams et al. (U.S. Pat. 5,550,318) and Collins et al. (U.S. Pat. 5,024,944). Methods for transforming wheat with such genes are described in Fry et al. (U.S. Pat. 5,631,152).

Other Embodiments

The invention further includes the use of analogs of any naturally-occurring MAPKKK polypeptide, a MAPKK polypeptide, or a WRKY polypeptide. Analogs
5 can differ from the naturally-occurring kinase domain by amino acid sequence differences, by post-translational modifications, or by both. Analogs of the invention will generally exhibit at least 40%, more preferably 50%, and most preferably 60% or even having 70%, 80%, or 90% identity with all or part of a naturally-occurring kinase domain amino acid sequence. The length of sequence
10 comparison is at least 15 amino acid residues, preferably at least 25 amino acid residues, and more preferably more than 35 amino acid residues. Modifications include in vivo and in vitro chemical derivatization of polypeptides, e.g., acetylation, carboxylation, phosphorylation, or glycosylation; such modifications may occur during polypeptide synthesis or processing or following treatment with isolated
15 modifying enzymes. Analogs can also differ from the naturally-occurring kinase domain polypeptide by alterations in primary sequence. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethyl methylsulfate or by site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory*
20 *Manual* (2d ed.), CSH Press, 1989, or Ausubel et al., *supra*). Also included are cyclized peptides, molecules, and analogs which contain residues other than L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g., or amino acids.

In addition, the invention also includes kinase domain fragments. As used
25 herein, the term "fragment," means at least 50 contiguous amino acids, preferably at least 130 contiguous amino acids, more preferably at least 160 contiguous amino acids, and most preferably at least 190 to 230 or more contiguous amino acids. Fragments of kinase domain polypeptides can be generated by methods known to those skilled in the art or may result from normal protein processing (e.g., removal
30 of amino acids from the nascent polypeptide that are not required for biological activity or removal of amino acids by alternative mRNA splicing or alternative protein processing events). In preferred embodiments, a kinase domain fragment (e.g., a fragment of NPK1, ANP1, ANP2, or ANP3) is capable of activating the

transcription of a plant defense gene. Methods for evaluating such activity are described herein.

All publications and patent applications mentioned in this specification are herein incorporated by reference.

5 Other embodiments are with the claims.

What is claimed is:

Claims

1. A method of enhancing resistance to a plant pathogen in a plant, said method comprising:

5 (a) providing a transgenic plant cell that expresses an isolated DNA molecule encoding a kinase domain of a MAPKK polypeptide; and

(b) regenerating a plant from said plant cell wherein said isolated DNA molecule is expressed in said plant, and wherein said plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant.

10

2. The method of claim 1, wherein said plant is a dicot.

3. The method of claim 2, wherein said dicot is a crucifer.

15

4. The method of claim 3, wherein said crucifer is *Arabidopsis*.

5. The method of claim 1, wherein said plant is a monocot.

20

6. The method of claim 1, wherein said kinase domain is constitutively active.

7. The method of claim 1, wherein said MAPKK polypeptide is MKK4.

8. The method of claim 1, wherein said MAPKK polypeptide is MKK5.

25

9. The method of claim 1, wherein said MAPKK polypeptide activates a gene involved in pathogen defense.

30

10. The method of claim 1, wherein said MAPKK polypeptide activates the PAL1, GST1, WRKY29, or PR1 gene promoters.

11. A method of enhancing resistance to a plant pathogen in a plant, said method comprising:

(a) providing a plant cell that expresses an isolated DNA molecule encoding a kinase domain of a MAPKKK polypeptide; and

5 (b) regenerating a plant from said plant cell wherein said isolated DNA molecule is expressed in said plant, and wherein said plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant.

12. The method of claim 11, wherein said plant is a dicot.

10

13. The method of claim 12, wherein said dicot is a crucifer.

14. The method of claim 13, wherein said crucifer is *Arabidopsis*.

15

15. The method of claim 11, wherein said plant is a monocot.

16. The method of claim 11, wherein said kinase domain is constitutively active.

20

17. The method of claim 11, wherein said MAPKKK polypeptide is MEKK1.

18. The method of claim 11, wherein said MAPKKK polypeptide is ANP1.

25

19. The method of claim 11, wherein said MAPKKK polypeptide activates a gene involved in pathogen defense.

20. The method of claim 11, wherein said MAPKKK polypeptide activates the PAL1, GST1, WRKY29, or PR1 gene promoters.

30

21. A method of enhancing resistance to a plant pathogen in a plant, said method comprising the steps of:

(a) providing a plant cell that expresses an isolated DNA molecule encoding a polypeptide comprising a polypeptide having substantial identity to a WRKY polypeptide; and

(b) regenerating a plant from said plant cell wherein said isolated DNA molecule is expressed in said plant, and wherein said plant has enhanced resistance to a plant pathogen compared to a corresponding untransformed plant.

22. The method of claim 21, wherein said plant is a dicot.

23. The method of claim 22, wherein said dicot is a crucifer.

24. The method of claim 23, wherein said crucifer is *Arabidopsis*.

25. The method of claim 21, wherein said plant is a monocot.

26. The method of claim 21, wherein said WRKY polypeptide induces its own expression.

27. An isolated nucleic acid molecule having a nucleotide sequence for a promoter that is capable of initiating pathogen-inducible transcription in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:

a) a nucleotide sequence comprising the sequence set forth in Figures 15 or 16;

b) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in Figures 15 or 16; and

c) a nucleotide sequence that has at least about 70% sequence identity to a sequence set forth in a) or b).

28. A DNA construct comprising a nucleotide sequence of claim 27 operably linked to a heterologous nucleotide sequence of interest.

29. A vector comprising the DNA construct of claim 27.

30. A host cell having stably incorporated in its genome the DNA construct of claim 27.

5

31. A method for expressing a heterologous nucleotide sequence in a plant, said method comprising transforming a plant cell with a DNA construct comprising said heterologous nucleotide sequence operably linked to a promoter that is capable of initiating transcription in a plant cell and regenerating a stably transformed plant from said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

- 10
- a) a nucleotide sequence comprising the sequence set forth in Figures 15 or 16;
 - b) a nucleotide sequence comprising at least 40 contiguous nucleotides of the
 - 15 sequence set forth in Figures 15 or 16; and
 - c) a nucleotide sequence that has at least about 70% sequence identity to a sequence set forth in a) or b).

20 32. The method of claim 31, wherein said plant is a dicot.

33. The method of claim 32, wherein said dicot is a crucifer.

34. The method of claim 33, wherein said crucifer is *Arabidopsis*.

25 35. The method of claim 31, wherein said plant is a monocot.

36. A plant cell stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating transcription in said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

- 5 a) a nucleotide sequence comprising the sequence set forth in Figures 15 or 16;
- b) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in Figures 15 or 16; and
- c) a nucleotide sequence that has at least about 70% sequence identity to a
- 10 sequence set forth in a) or b).

37 The plant of claim 36, wherein said plant is a dicot.

38. The plant of claim 37, wherein said dicot is a crucifer.

15

39. The plant of claim 28, wherein said crucifer is *Arabidopsis*.

40. The plant of claim 36, wherein said plant is a monocot.

20

41. Seed of the plant of any one of claims 36-40.

A MAPK Signaling Pathway Leading To WRKY29 Induction





FIGURE 1

WRKY29



Target Sequence : W box
(T)(T)TGAC(C/T)

FIGURE 2

Flg22 Induces WRKY29 in Arabidopsis Protoplasts (RT-PCR Analysis)




FIGURE 3

MRK4 (cDNA accession number: AB015315)

Wild-type cDNA sequence from to end codons (1101 base pairs including the end TAG):

[illegible]

Wild-type protein sequence (366 aminoacids):

MRPIQSPGVSPVKSRPRRBDLTPLPQDVSLSLAVPLPLPTSGSGSGSAPS
SGSASSTNTNSSIEAKNYSDLVRGNRI GSGAGGTVYKVIHRPSSRLYATLKVIYGNH
EETVRQICREIETRDVNHPNVKCHMFQDNGEIQVLTFFMDKGSLEGAHVMEQ
QIADLSRQILSGALYLSHRHIVHRDIKPSNLTINSKXKIVADFGVSRILAQMDPC
NBSVGTIAYMSPERINTDLNQGKYDYGADIMSLGVSIIEFYLGFPFVPSRQGDWA
SLMCAICMSQPEAPATAPEFRHFISSCQLREBGRKSAMQLLQHPFITRASPSQN
RSPQNLHQLTLPFRPLSSSSSPLT (SEQ ID NO.:2)

FIGURE 4

Mutations rendering MKK4 constitutively active:

modify Serine (S) 230 to Glutamic Acid (E) by changing codon TCA into GAA

The mutations were done by PCR using the primers (mutated base pairs in lower case, both are from 5' end to 3' end):

CTTGGCTCAGTATGGATCCGTGTAATGAATCTGTTGGAAC
TCCAACAGATtcATTACACGGATCCATACTGAGCCAAG (SEQ ID NO.3)

So the sequences after mutations are:

**MKK4act mutant sequence from start to end codons
(1101 base pairs including the end TAG):**

agaccgattcaatcgctccaggagtttccggtccggtgaaaagccgtcccgctcgccgtcctgat
cttaccttacgcttcccaacgcgatgtttctctcgctgtacctcttccctctccacctaactccggt
ggttccggtggctctagtggatctcgccgctcttctgggtggttcggcgctcttcaacgaacactaacagc
tccatagaagcgaagaactattcggatttagtgagaggttaaccgatcgggaagcggagcaggtggaacg
gtatacaaagtgattcacgcgtccgagttctcgctatatgcacttaagggtgatatacggtaaccacgag
gagactgtgagacgtcagatctgtagagagatcgagattttacgagatgtgaatcatccaaacggttg
aaatgtcacgagatgtttgatcagaacggtgagatccagggttttgcttgagtttatggataaaggttct
ttagaagggtgctcatgtgtggaagagcaacaattagctgatctatctcgctcagattcttagtggttta
gcttatctccatagccgtcacatagttcatcgatgatatcaaacatcgaatcttttgataaactctgct
aaaaacggttaagattgctgattttggagtttagtaggatcttggtcagtatggatccgtgtaattgaa
tctgttgaaccattgcttatatgagtcctgagaggattaacactgatttgaatcagggaagtatgat
ggttatgctggagatatttggagcttaggtgttagcattttggagtttacttggggagggttcccttc
cctgtgagtagacaagggtgattgggctagtcttatgtgtgccatttgtatgtctcagcctccagaagct
ccagcgactgcgtcgccggagtttcggcattttatctcggtgttgcagagagaaccggggaaaagg
aggagtgcattgcagctattgcagcatcctttcatattaagagcaagtcggagccagaacaggtctcct
cagaatctacatcaactcttgccctcctcctcgctcctctgtcctcgtcttcttccaaccacatag
(SEQ ID NO.:4)

MKK4act mutant protein sequence (366 aminoacids):

MRPIQSPPGVSVPVKSRPRRRPDLTLPLPQRDVSLAVPLPLPPTSGGSGGSSGSAPSSGGSASSTNTNS
SIEAKNYSIDLVRGNRIGSGAGGTVYKVIHRPSSRLYALKVIYGNHEETVRRQICREIEILRDVNHPNVV
KCHEMFDQNGEIQVLLFMDKGSLEGAVWKEQQLADLSRQILSGLAYLHSRHIVHRDIKPSNLLINSA
KNVKIADFGVSRILAQMDPCNEISVGTIAYMSPERINTDLNQKYGAGDIWSLGVSILEFYLGFRFPF
PVSROGDWASLMCAICMSQPPEAPATASPEFRHFISCCLOREPFGKRRSAMQLLOHPFILRASPSQNRSP
QNLHQLLPPPRPLSSSSSPTT (SEQ ID NO.:5)

FIGURE 5

MKK5 (cDNA accession number: AB015316)

Wild-type cDNA sequence from Strain to end codons (1047 base pairs including the end TAG):

[illegible]

Wild-type protein sequence (348 aminoacids):

MKPIQSPSGVAPBMKNRLRKRPDLSTPLPHRDVATLAVPLPBPSSSSAPASSAI
 STNISAAKSLSETERVNRIGSGAGGLVYKVIHTLTSRPFATKVIYGNHEDTVRRQIC
 REIEITRVSVDHPNVKCHDMFDHNGEIQVLEFMDGSLGSAHIMQEQLEIADLSROI
 LSGTAYLHRHIVHRDIKPSNLTINSKXKIVADFGVSRILAQMDPCNSVGTIAY
 MSPERINTDLNHGRYDAGDVMSLGVSILEYILGRFPFAVSRQGDWASIMCAICMS
 QPPEAPATASQEFFRHFVSCCIQSDPPKRWMSAQQLTQHPFIILKATGGPNLROMLPBR
 PLPSAS (SEQ ID NO.: 7)

FIGURE 6

Mutations rendering MKK5 constitutively active:

modify Serine (S) 221 to Glutamic Acid (E) by changing codon TCA into GAA

The mutations were done by PCR using the primers (mutated base pairs in lower case, both are from 5' end to 3' end):

CTTGGCACAA**gaa**AATGGATCCTTGTAAT**ga**ATCTGTTGGT
ACCAACAGAT**tc**CATTACAAGGATCCATT**tt**TGTGCCAAG (SEQ ID NO.:8)

So the sequences after mutations are:

MKK5act mutant cDNA sequence from **552 to **end** codons**

(1047 base pairs including the end TAG):

aaaccgattcaatctccttctggagtagcttcacctatgaagaaccggtttacgc
aaacgtcctgacctaaagcttaccactcccacaccgcgacgtcgctctcgccgtacct
ctccctctcccacctccttcttctcttcatccgctccggcgctcttctccgcgcatc
tcaaccaacatctccgccgctaaaagcttatccgagctagaacgagtgaaacgaatc
ggaagcggagccggaggaacggtttacaaagtaatccacactccgacgtcacgtcct
ttcgctctcaaagtgatttacggaaaccacgaagataccgtgagacgtcagatctgt
agagagatcgagatcttaagaagtgttgatcatcaaacggttgtaaattgtcacgat
atgtttgatcataacggtgagatccaggttttgcttgagtttatggatcaaggatct
cttgaaggagctcatatatggcaagaacaggaattagctgatctctctcgctcagatt
cttagtgattagcttatcttcatcgctcgctcatatcgttcatcgatgatcaaacct
tcgaatctccttataaaactcagctaaaaatgtgaaaattgctgattttggtgtgagt
aggatcttggcacaa**ga**aatggatccttgtaat**ga**atctgttggtactattgcttat
atgagtcctgagaggattaatactgatttgaatcatggctcggttacgatggttatgct
ggagatgtttggagttaggtgttagtatcttggagttttacttggggaggtttcct
tttgctgtgagtagacaaggtgattgggctagtcttatgtgtgctatttgatgtct
cagccacctgaagctccggctacggcgctcaggagtttcgtcactttgtttcttgt
tgtttacagagtgatecctcctaagagatggctcagctcaacagcttttgcagcatcct
ttcataacttaaagctaccggtggtcctaatactccgtcaaagtgtgccgccgctcgt
cctcttcttctgcctcttag (SEQ ID NO.:9)

MKK5act mutant protein sequence (348 aminoacids):

MKPIQSPSGVASPMKNRLRKPDL~~SL~~PLPHRDVALAVPLPLPPSSSSSSAPASSSAI
STNISAAKSLSELERVNRIGSGAGGT~~VY~~KVIHTPTSRPFALKVIYGNHEDTVRRQIC
REIEILRSVDHPNVVKCHDMFDHNGEIQVLLFMDQGSLEGAHIWQEQLADLSRQI
LSGLAYLHRRHIVHRDIKPSNLLINSAKNVKIADFGVSRILAQ**MDPCN**ESVGTIAY
MSPERINTDLNHGRYDGYAGDVWSLGV~~S~~ILEFYLG~~R~~FPFAVSROGDWASLMCAICMS
QPPEAPATASQEFRHFVSCCLQSDPPKRWSAQQLLQHPFILKATGGPNL~~RQ~~MLPPPR
PLPSAS (SEQ ID NO.:10)

FIGURE 7

Alignment of MKK4 and MKK5 wild-type:


```

MKK4 MRPIQSPGVSPVKSRPRRPDLTLPLPQRDVSLAVPLPLPTSGSGG
MKK5 MKPIQSPGVASPMKNRLTKRPDLTLPLPHRDVALAVPLPLPPS-----
MKK4 SSGAPSSGSGASSTNTNSSIBAKNYSDLVRGNRIGSGAGGVYKVIHRP
MKK5 SSSAPASS-SAISTNIS--AAKSLSELERVNRRIGSGAGGVYKVIHTP
MKK4 SSRLYATKVIYGNHHEETVRRQICREIIEILRDVNHHPNVVKCHDMFDNGEI
MKK5 TSRPFATKVIYGNHHEDTVRRQICREIIEILRSVDHPNVVKCHDMFDHNGEI
MKK4 QVLTFMDKGSLEGAWKKEQQLADLSRQILSGLAYLHSHRHIVHRDIKPS
MKK5 QVLTFMDQGSLEGAHIWQEQELADLSRQILSGLAYLHRRHIVHRDIKPS
MKK4 NLTINSAKNVKIADFGVSRILAQIMDPGNSVGTIAYMSPERINTDINH
MKK5 NLTINSAKNVKIADFGVSRILAQIMDPGNSVGTIAYMSPERINTDINH
MKK4 KYDGYAGDIWSLGVSIIEFYILGRFPFVSRQGDWASLMCAICMSQPPBAP
MKK5 RYDGYAGDVWSLGVSIIEFYILGRFPFVSRQGDWASLMCAICMSQPPBAP
MKK4 ATASPEERHFIISCCLOREBPGRKRSAMQTLQHPFIIRASPQNRLHQ
MKK5 ATASQEFFRHFFVSCCLQSDPPKRWSAQQLQHPFIILKATG----GP-NLRQ
MKK4 LTPPPRPLSSSSSPTT
MKK5 MLPPRPPLPSAS-----
      (SEQ ID NO.: 11)
      (SEQ ID NO.: 12)

```

FIGURE 8

Constitutively Active AtMEKK1 Induces WRKY29



GFP MEKK1 CTR1




FIGURE 9

Constitutively Active AtMKK4 Induces WRKY29





FIGURE 10

WRKY29 Induces Its Own Promoter




Fig22




FIGURE 11

WRKY29 Regulates Early Defense Genes





FIGURE 12

Agrobacterium-Mediated Transient Transfection




FIGURE 13

Transient Expression of WRKY29 Reduces Pathogen Susceptibility




FIGURE 14





WRKY29 promoter fragment

5'-


TCGTCCATGATGGACATATATGTGACTGCGTATTTACACACACCGCACGTATGCTTATTTTACACA
CGTTAGAAGAAGAATTCAAAGAAGTCGGTTCTATTTGTATTCTTGTGAGATCATCAATATGAC
AATATCGTTCTATTAATATACGTATAATTCATATGTTGTCATGGTTTCACATACCATGTCGACA
GTCGACGTACGTACAAAAGTATAAATAGTATGAATCTAATAACAGCACCAAGATTGAAGTTC
ATCTTCTAATCAAACTATCATAAAGTGGTTTCAAAATAGTGTTTTTCTGATGAAACTATAAC
TGAGTTATAATCAATCCGAAATTATATACTAATTATATTTGGGAACTAGATAAACGCAAAAA
CATGAGCAGTTTCTTATTTTTTTGTCCAGATTTAAAAATTGGAGTGTTAAAAATATACGGAGTG
TTACACAATGAAAACACAAGAAGTCAAGAACCCATAAGTTATTTTAATTAATAATATTGTATT
TAAAGTGATTATTAAAAAATAATGTAAAACTGATTATTTGTTGACAAAAAACAGTTAGTTAT
AGTTAAATAGTATTGATGCATATATATACTATCTCATTATTTTGGTATTACTCAGTACTCACAT
CTTTAATAAAGACAAAGATAGTTAGTGTATAATTCAAATCGAACTCACAGAAGTCAATAAGC
GCGTAAAAATACAAAATATCTGGCAGACTTTAGCAAGGTTTGTTCACAGAAATGGTCA
TTTCAGAATCATTTACATATCCATATATATAGCTCTTAAATGGTATATATTGGGTAATGGGTAT
TCGTTTAAATAATTTGTTTTCTGTAAATTTCAAATATTAATCTGATCAGTTTATCCATGTGTGT
ATATTTAGTGTATTATCATCAATATATGACATAGACAGACTTTCAAGTTGGTGCAAGAGGGGA
TGAAAATTTCTTCCAGTTGCAAGAGTAAGCTGACTAGCATTTTTTTTTATATATAATTTATTT
CTCAAATGGTTTTTATTATTGTTTTGTTGACTTTAAAGTTTTTGCTTTTATGGGACTGCAATCAC
CCGTGCCAATTTTAAATCTCCATCGCCTAAAAAAGAAAGAAAAGGCTACCATTATGGACCGAA
ATATTTAAGACCATAATACAAAATTATACGAATATTTTCTGTAATTATATATATGATCATTG
ACAAAGCAAAGTCAATCAAATAAACTTCAAAGAAATTATGAGCTTATAATAAGTTTGATAGT
GTTAAATATAATCAAATCATTAATTTAGTATTTATTTTCATCTCGGTTTCCATTTGATAGATAG
ATAAATGAAAGATAGCATCGCCAATAATGAAAAACTTTATTTGATGGCAATACTTTGTTACA
TCATTTCTGTTTTCTTAATTTTCATGTCGAAATATTGCCATGATTGTGTTCAACATAACTAGTTTT
GAGGTAACAAGTTAAAAATTTGTTATATTTTTGAATATGTTATTCAGTTGAAAGTCATTTAGA
TGTAAGTAAAAACAAACATAAGAAGTTAACATATCAATATTAAACACAGCGAATAATCATTATT
ACAAAAAAAAGCAAAAAAATAGAAGAAGATATTATATATTGGAGAATCTTTCTTTAGTCTTA
GTTGGGAAGATTTTGTGCTATGGGATTAAAGGTATCCATCCCTATTCTATGATAGAGGCGTGG
GGTTATTGGACCAATCTATATATATTACCACAAGGCTTAAGATGAAGTGATAATACAGTATTA
TTAATACCCTCCCAAATTATTTTTAAATATTTATCAAAAGAAGCTTACGGTATAGATCATACTT
GCAGCATTATTCTATAAGTTTATTTAATTTTCAGTGGCTCGTTACGTGAACACAAGGTAAGCTA
ATAGACTTACGTGCCCCATTAAACACATACATAATTATACAAGTATCATGAACTAGTGACAA
AACCTCGATCAAATAAAGAAATTACCATGACGACAAAAGATAATTAAAAAAAACACTACTAT
ATGTCATACTCATGCATATGCATGTACAAATGCCGCTTTAAATATTTAATTTAGTTAAAGCAAT
GATATTTAAATTTCTCTCTACTTCATATATATTTCCAAAAGACATATTGTCAAATTCCTTTTTT
TTATATATAATCATATATTCATATTGTTATATTTTCAATATTTAATAGTAAGATGGACTTTCCTG
AATGTTGTGTATGATTTATAATTTGAGATATTTTGTGCGGAGATGGATATTTGACAAGTTAATGT
TACTTTATTAATAATTTCTAAACATTTAGGTACGAATTGACTTTTTTCAAAGTCAACACAATAA
ATTTTAAAGTTTAATGACTTAACGGGTTACATGGGAAACGAAAACACCCTAAACCACAAA
CAATCTAATCTTATTTCTTTTATATAAACCGCTGTTTCCAAAAGGCTTGTTCCTCGTCATAT
GTACTTGTACACCAACCCACCAAAAGAGATAAAAGAGGAAACAAAACTCGAAAAGAGAGA
GATATATGGGTGAGGTGGCTTAT - 3' (SEQ ID NO.: 13)

Figure 15


Figure 16



FIGURES 17 A-D




FIGURES 18 A-D




FIGURES 19 A-E


20 A




20 B




20 C




20 D





FIGURES 21 A-F

FIGURE 22



SEQUENCE LISTING

<110> The General Hospital Corporation

<120> Master Activators of Pathogen Responsive Genes

<130> 00786/397WO2

<150> US 60/275,199

<151> 2001-03-12

<160> 14

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1101

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```

atgagaccga ttcaatcgcc tccaggagtt tccgttccgg tgaaaagccg tcccgcgcgc 60
cgtcctgata ttaccttacc gcttcctcaa cgcgatgttt ctctcgctgt acctcttctc 120
ctcccaccta ctcccggtgg ttccggtggc tctagtggat ctgcccgcgc ttctgggtgg 180
tcggcgctctt caacgaacac taacagctcc atagaagcga agaactattc ggatttagtg 240
agaggtaacc gtatcggaag cggagcaggt ggaacgggat acaaagtgat tcaccgtccg 300
agttctcgtc tatatgcact taaggtgata tacggtaacc acgaggagac tgtgagacgt 360
cagatctgta gagagatcga gattttacga gatgtgaatc atccaaacgt tgtgaaatgt 420
cacgagatgt ttgatcagaa cggtgagatc caggttttgc ttgagtttat ggataaagg 480
tctttagaag gtgctcatgt gtggaaagag caacaattag ctgatctatc tcgtcagatt 540
cttagtggtt tagcttatct ccatagccgt cacatagtcc atcgtgatat caaaccatcg 600
aatcttttga taaactctgc taaaaacggt aagattgctg attttgagg attttgagg 660
ttggctcaga ctatggatcc gtgtaattca tctgttgga ccattgctta tatgagtcct 720
gagaggatta acactgattt gaatcagggg aagtatgatg gttatgctgg agatatttgg 780
agcttaggtg ttagcatttt ggagttttac ttggggaggt ttcctttccc tgtgagtaga 840
caaggtgatt gggctagtct tatgtgtgcc atttgatgt ctcagcctcc agaagctcca 900
gcgactgcgt cgccggagtt tcggcatttt atctcgtgtt gcttgagag agaaccgggg 960
aaaaggagga gtgctatgca gctattgcag catcctttca tattaagagc aagtcagagc 1020
cagaacaggt ctctcagaa tctacatcaa ctcttgctc ctctcgtcc tctgtcctcg 1080
tcttcttctc caaccacata g

```

<210> 2

<211> 366

<212> PRT

<213> Arabidopsis thaliana

<400> 2

```

Met Arg Pro Ile Gln Ser Pro Pro Gly Val Ser Val Pro Val Lys Ser
1           5           10           15
Arg Pro Arg Arg Arg Pro Asp Leu Thr Leu Pro Leu Pro Gln Arg Asp
20           25           30
Val Ser Leu Ala Val Pro Leu Pro Leu Pro Pro Thr Ser Gly Gly Ser
35           40           45
Gly Gly Ser Ser Gly Ser Ala Pro Ser Ser Gly Gly Ser Ala Ser Ser
50           55           60
Thr Asn Thr Asn Ser Ser Ile Glu Ala Lys Asn Tyr Ser Asp Leu Val

```

65 Arg Gly Asn Arg Ile Gly Ser Gly Ala Gly Thr Val Tyr Lys Val 85
 70 Ile His Arg Pro Ser Ser Arg Leu Tyr Ala Leu Lys Val Ile Tyr Gly 90
 75 Asn His Gln Thr Val Arg Arg Gln Ile Cys Arg Gln Ile Gln Ile 105
 110 Leu Arg Asp Val Asn His Pro Asn Val Val Lys Cys His Gln Met Phe 115
 120 Leu Arg Asp Val Asn His Pro Asn Val Val Lys Cys His Gln Met Phe 125
 130 Asp Gln Asn Gly Gln Ile Gln Val Leu Leu Gln Phe Met Asp Lys Gly 135
 140 Ser Leu Gln Gly Ala His Val Trp Lys Gln Gln Leu Ala Asp Leu 145
 150 Ser Arg Gln Ile Leu Ser Gly Leu Ala Tyr Leu His Ser Arg His Ile 155
 160 Val His Arg Asp Ile Lys Pro Ser Asn Leu Leu Ile Asn Ser Ala Lys 165
 170 Asn Val Lys Ile Ala Asp Phe Gly Val Ser Arg Ile Leu Ala Gln Thr 175
 180 Met Asp Pro Cys Asn Ser Ser Val Gly Thr Ile Ala Tyr Met Ser Pro 185
 190 Gln Arg Ile Asn Thr Asp Leu Asn Gln Gly Lys Tyr Asp Gly Tyr Ala 195
 200 Gly Asp Ile Trp Ser Leu Gly Val Ser Ile Leu Gln Phe Tyr Leu Gly 205
 210 Arg Phe Pro Phe Val Ser Arg Gln Gly Asp Trp Ala Ser Leu Met 215
 220 Cys Ala Ile Cys Met Ser Gln Pro Gln Ala Pro Ala Thr Ala Ser 225
 230 Pro Gln Phe Arg His Phe Ile Ser Cys Cys Leu Gln Arg Gln Pro Gly 235
 240 Lys Arg Arg Ser Ala Met Gln Leu Leu Gln His Pro Phe Ile Leu Arg 245
 250 Ala Ser Pro Ser Gln Asn Arg Ser Pro Gln Asn Leu His Gln Leu 255
 260 Pro Pro Arg Pro Leu Ser Ser Ser Ser Ser Pro Thr Thr 265
 270 Ala Ser Pro Phe Pro Val Ser Arg Gln Gly Asp Trp Ala Ser Leu Met 275
 280 Arg Phe Pro Phe Pro Val Ser Arg Gln Gly Asp Trp Ala Ser Leu Met 285
 290 Cys Ala Ile Cys Met Ser Gln Pro Gln Ala Pro Ala Thr Ala Ser 295
 300 Pro Gln Phe Arg His Phe Ile Ser Cys Cys Leu Gln Arg Gln Pro Gly 305
 310 Lys Arg Arg Ser Ala Met Gln Leu Leu Gln His Pro Phe Ile Leu Arg 315
 320 Ala Ser Pro Ser Gln Asn Arg Ser Pro Gln Asn Leu His Gln Leu 325
 330 Ala Ser Pro Ser Gln Asn Arg Ser Pro Gln Asn Leu His Gln Leu 335
 340 Pro Pro Arg Pro Leu Ser Ser Ser Ser Ser Pro Thr Thr 345
 355

<210> 3
 <211> 82
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 3
 cttgctcag gatatgagtc cgtgtaatga atcgtgtgga actccaacag attcattaca 60
 cggatccata tccctgagcca ag
 <210> 4
 <211> 1101
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 4
 atgagaccga ttcaatcgcc tccaggagtt tccgttcggg tgaagaagcgg tcccggtcgc 60
 cgtccctgac ttaccctacc gcttccctcaa cgcgatgttt cttcgcgtgt accctctcct 120
 ctcccaacct cttccggtgg tccggtgggc tctagtggat ctgcgcgcgtc ttctggtgtg 180
 tcggcgctct caacgaacac taacagctcc atagaagcga agaactatc ggatttagtg 240


```

agaggttaacc gtatcggaag cggagcaggt ggaacgggtat acaaagtgat tcaccgtccg 300
agttctcgtc tatatgcact taaggtagata tacggtaacc acgaggagac tgtgagacgt 360
cagatctgta gagagatcga gattttacga gatgtgaatc atccaaacgt tgtgaaatgt 420
cacgagatgt ttgatcagaa cggtagatc caggttttgc ttgagttat ggataaagggt 480
tctttagaag gtgctcatgt gtggaaagag caacaattag ctgatctatc tcgtcagatt 540
cttagtggtt tagcttatct ccatagccgt cacatagttc atcgtgatat caaacatcg 600
aatcttttga taaactctgc taaaaacggt aagattgctg attttggagt tagtaggac 660
ttggctcagg atatggatcc gtgtaatgaa tctgttgga ccatgctta tatgagtcct 720
gagaggatta acactgattt gaatcaggga aagtatgatg gttatgctgg agatatttgg 780
agcttaggtg ttagcatttt ggagttttac ttggggaggt ttcttttccc tgtgagtaga 840
caaggtgatt gggctagtct tatgtgtgcc atttgatgt ctcagcctcc agaagctcca 900
gcgactcgct cgccggagtt tcggcatttt atctcgtgtt gcttcagag agaaccgggg 960
aaaaggagga gtgctatgca gctattgcag catcctttca tattaagagc aagtcggagc 1020
cagaacaggt ctctcagaa tctacatcaa ctcttgctc ctctcgtcc tctgtcctcg 1080
tcttcttctc caaccacata g 1101

```

<210> 5

<211> 366

<212> PRT

<213> Arabidopsis thaliana

<400> 5

```

Met Arg Pro Ile Gln Ser Pro Pro Gly Val Ser Val Pro Val Lys Ser
1      5      10      15
Arg Pro Arg Arg Arg Pro Asp Leu Thr Leu Pro Leu Pro Gln Arg Asp
20      25      30
Val Ser Leu Ala Val Pro Leu Pro Leu Pro Pro Thr Ser Gly Gly Ser
35      40      45
Gly Gly Ser Ser Gly Ser Ala Pro Ser Ser Gly Gly Ser Ala Ser Ser
50      55      60
Thr Asn Thr Asn Ser Ser Ile Glu Ala Lys Asn Tyr Ser Asp Leu Val
65      70      75      80
Arg Gly Asn Arg Ile Gly Ser Gly Ala Gly Gly Thr Val Tyr Lys Val
85      90      95
Ile His Arg Pro Ser Ser Arg Leu Tyr Ala Leu Lys Val Ile Tyr Gly
100     105     110
Asn His Glu Glu Thr Val Arg Arg Gln Ile Cys Arg Glu Ile Glu Ile
115     120     125
Leu Arg Asp Val Asn His Pro Asn Val Val Lys Cys His Glu Met Phe
130     135     140
Asp Gln Asn Gly Glu Ile Gln Val Leu Leu Glu Phe Met Asp Lys Gly
145     150     155     160
Ser Leu Glu Gly Ala His Val Trp Lys Glu Gln Gln Leu Ala Asp Leu
165     170     175
Ser Arg Gln Ile Leu Ser Gly Leu Ala Tyr Leu His Ser Arg His Ile
180     185     190
Val His Arg Asp Ile Lys Pro Ser Asn Leu Leu Ile Asn Ser Ala Lys
195     200     205
Asn Val Lys Ile Ala Asp Phe Gly Val Ser Arg Ile Leu Ala Gln Asp
210     215     220
Met Asp Pro Cys Asn Glu Ser Val Gly Thr Ile Ala Tyr Met Ser Pro
225     230     235     240
Glu Arg Ile Asn Thr Asp Leu Asn Gln Gly Lys Tyr Asp Gly Tyr Ala
245     250     255
Gly Asp Ile Trp Ser Leu Gly Val Ser Ile Leu Glu Phe Tyr Leu Gly
260     265     270
Arg Phe Pro Phe Pro Val Ser Arg Gln Gly Asp Trp Ala Ser Leu Met
275     280     285

```

Cys Ala Ile Cys Met Ser Gln Pro Pro Gln Ala Pro Ala Thr Ala Ser
 290 295 300 310 315 320 325 330 335 340 345 350 355 360 365
 Pro Gln Phe Arg His Phe Ile Ser Cys Cys Leu Gln Arg Gln Pro Thr Thr
 Ala Ser Pro Ser Gln Asn Arg Ser Pro Gln Asn Leu His Gln Leu Leu
 Pro Pro Pro Arg Pro Leu Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser

<210> 6
 <211> 1047
 <212> DNA

<213> Arabidopsis thaliana

atgaaaccca ttcaatccc ttctggagta gctcaccta ttgaagaaccg tttaacgcaaa 60
 cgtccctgacc taagcttacc actccacac ccggaagtcg ctctcgccgt acctccct 120
 ctccaccctc ctctctccct ttcatccgct ccggcgctct cctccgcgat ctcaaccaac 180
 atctccgcgcg ctaaaagctt atccggagcta gaacggagtg accgaatcgg aagcggaacc 240
 ggaaggaaacgg tttaacaagt aatccacac ccgaagtcac gtcctttcgc tctcaaatg 300
 attaacggaa accaaggaaga taccgtagaga cgtcagatct gtagagagat cgagatctta 360
 agaaagtggtt atcatccaaa cgttgtagaaa tgytcaagata tgyttagata taacggtagag 420
 atccagggtt tgcctgagta taggatacga ggaatccttg aaggagctca tatacgcaaa 480
 gaacaggaaat tagctgatat ctctcgtag atctcttagtg aaggaagctca tatacgcaaa 540
 cgtcatatcg ttcatcagtg tatcaaaact tcgaatccc ttataaactc agctaaaat 600
 gtgaaaaatg ctgattttgg tgtgagtagg atcttgagca aacaatgga tccctgtaat 660
 tcatctgttg gtaatatgc ttatatagtg cctgagagga ttaatactga ttgtaatat 720
 ggtcgttaag atggtatagc tggagatgtt tggagtttag ggtttagtat cttgaggtt 780
 tacttgggga ggttccctt tgcgtgtagt agacaagggtg atcgggctag tctatgtgt 840
 gctattgta gttcagacc acctgaagct ccgctaggg cgtctcagga gttcgtcac 900
 ttgtttctt ggtgttaca gagtatcct cctaagagat ggtcagctca acagctttg 960
 cagcatcctt tcatactaa agctaacggg ggtcccaatc tccgtcaaat gttgcccgcg 1020
 cctcgctcctc ttctctcgc ctcttag

<210> 7

<211> 348

<212> PRT

<213> Arabidopsis thaliana

<400> 7
 Met Lys Pro Ile Gln Ser Pro Ser Gly Val Ala Ser Pro Met Lys Asn
 1 5 10 15
 Arg Leu Arg Lys Arg Pro Asp Leu Ser Leu Pro Leu Pro His Arg Asp
 20 25 30
 Val Ala Leu Ala Val Pro Leu Pro Leu Pro Pro Ser Ser Ser Ser
 35 40 45
 Ser Ala Pro Ala Ser Ser Ala Ile Ser Thr Asn Ile Ser Ala Ala
 50 55 60
 Lys Ser Leu Ser Gln Leu Arg Val Asn Arg Ile Gly Ser Gly Ala
 65 70 75 80
 Gly Gly Thr Val Tyr Lys Val Ile His Thr Pro Thr Ser Arg Pro Phe
 85 90 95
 Ala Leu Lys Val Ile Tyr Gly Asn His Gln Asp Thr Val Arg Arg Gln
 100 105 110
 Ile Cys Arg Gln Ile Gln Ile Leu Arg Ser Val Asp His Pro Asn Val

115	120	125
Val Lys Cys His Asp Met Phe Asp His Asn Gly Glu Ile Gln Val Leu		
130	135	140
Leu Glu Phe Met Asp Gln Gly Ser Leu Glu Gly Ala His Ile Trp Gln		
145	150	155
Glu Gln Glu Leu Ala Asp Leu Ser Arg Gln Ile Leu Ser Gly Leu Ala		
165	170	175
Tyr Leu His Arg Arg His Ile Val His Arg Asp Ile Lys Pro Ser Asn		
180	185	190
Leu Leu Ile Asn Ser Ala Lys Asn Val Lys Ile Ala Asp Phe Gly Val		
195	200	205
Ser Arg Ile Leu Ala Gln Thr Met Asp Pro Cys Asn Ser Ser Val Gly		
210	215	220
Thr Ile Ala Tyr Met Ser Pro Glu Arg Ile Asn Thr Asp Leu Asn His		
225	230	235
Gly Arg Tyr Asp Gly Tyr Ala Gly Asp Val Trp Ser Leu Gly Val Ser		
245	250	255
Ile Leu Glu Phe Tyr Leu Gly Arg Phe Pro Phe Ala Val Ser Arg Gln		
260	265	270
Gly Asp Trp Ala Ser Leu Met Cys Ala Ile Cys Met Ser Gln Pro Pro		
275	280	285
Glu Ala Pro Ala Thr Ala Ser Gln Glu Phe Arg His Phe Val Ser Cys		
290	295	300
Cys Leu Gln Ser Asp Pro Pro Lys Arg Trp Ser Ala Gln Gln Leu Leu		
305	310	315
Gln His Pro Phe Ile Leu Lys Ala Thr Gly Gly Pro Asn Leu Arg Gln		
325	330	335
Met Leu Pro Pro Pro Arg Pro Leu Pro Ser Ala Ser		
340	345	

<210> 8
 <211> 80
 <212> DNA
 <213> Arabidopsis thaliana

<400> 8
 cttggcaca gaaatggatc cttgtaatga atctgttggt accaacagat tcattacaag 60
 gatccatttc ttgtgccaaag 80

<210> 9
 <211> 1047
 <212> DNA
 <213> Arabidopsis thaliana

<400> 9
 atgaaaccga ttcaatctcc ttctggagta gcttcaccta tgaagaaccg tttacgcaaa 60
 cgtcctgacc taagcttacc actccacac cgcgacgtcg ctctcgccgt acctctccct 120
 ctcccacctc cttcttcctc ttcatccgct cggcgctctt cctccgcat ctcaaccaac 180
 atctccgccg ctaaaagctt atccgagcta gaacgagtga accgaatcgg aagcggagcc 240
 ggaggaacgg tttacaaagt aatccacact ccgacgtcac gtcctttcgc tctcaaagtg 300
 atttacggaa accacgaaga taccgtgaga cgtcagatct gtagagagat cgagatctta 360
 agaagtgttg atcatccaaa cgttgtgaaa tgtcacgata tgtttgatca taacgggtgag 420
 atccagggtt tgcttgagtt tatggatcaa ggatctcttg aaggagctca tatatggcaa 480
 gaacaggaat tagctgatct ctctcgctcag attcttagtg gattagctta tcttcacgtg 540
 cgtcatatcg ttcacgtgta tatcaaacct tcgaatctcc ttataaactc agctaaaaat 600
 gtgaaaattg ctgattttgg tgtgagtagg atcttggcac aagaaatgga tccttgtaat 660
 gaatctgttg gtactattgc ttatatgagt cctgagagga ttaatactga tttgaatcat 720

ggtcgctacg atggttatgc tggagatgtt tggagtttag gtgttagtat cttagagttt 780
 tacttgaggga ggttcctt tgcgtgtagt agacaaggtg atgggctag tcttatgtgt 840
 gctatttgta tgcctcagcc acctgaagct ccggctcagcg cgtctcagga gtttcgtcac 900
 ttgtttctt gttggttaca gagtgcctt cctaagagat ggtcagctca acaagctttg 960
 cagcatcctt tcatacctaa agctaccggt ggtcctaat ccgccaat gttgcgcgcg 1020
 cctcgcttc ttcttctgc ctcttag

<210> 10
 <211> 348
 <212> PRT
 <213> Arabidopsis thaliana

Met Lys Pro Ile Gln Ser Pro Ser Gly Val Ala Ser Pro Met Lys Asn 1
 Lys Ser Leu Ser Gln Leu Gln Arg Val Asn Arg Ile Gly Ser Gly Ala 65
 Gly Gly Thr Val Tyr Lys Val Ile His Thr Pro Thr Ser Arg Pro Phe 80
 Ala Leu Lys Val Ile Tyr Gly Asn His Gln Asp Thr Val Arg Arg Gln 95
 Ile Cys Arg Gln Ile Gln Ile Leu Arg Ser Val Asp His Pro Asn Val 110
 Val Lys Cys His Asp Met Phe Asp His Asn Gly Gln Ile Gln Val Leu 125
 Leu Gln Phe Met Asp Gln Gly Ser Leu Gln Gly Ala His Ile Trp Gln 140
 Leu Gln Phe Met Asp Gln Gly Ser Leu Gln Ile Leu Ser Gly Leu Ala 155
 Tyr Leu His Arg Arg His Ile Val His Arg Asp Ile Lys Pro Ser Asn 170
 Ser Arg Ile Leu Ala Gln Gln Met Asp Pro Cys Asn Gln Ser Val Gly 205
 Thr Ile Ala Tyr Met Ser Pro Gln Arg Ile Asn Thr Asp Leu Asn His 220
 Gly Arg Tyr Asp Gly Tyr Ala Gly Asp Val Trp Ser Leu Gly Val Ser 235
 Ile Leu Gln Phe Tyr Leu Gly Arg Phe Pro Phe Ala Val Ser Arg Gln 255
 Gly Asp Trp Ala Tyr Asp Gly Tyr Ala Gly Asp Val Trp Ser Leu Gly Val 270
 Gly Asp Trp Ala Ser Leu Met Cys Ala Ile Cys Met Ser Gln Pro Pro 285
 Gln Ala Pro Ala Thr Ala Ser Gln Gln Phe Arg His Phe Val Ser Cys 300
 Cys Leu Gln Ser Asp Pro Lys Arg Trp Ser Ala Gln Gln Leu Leu 315
 Gln His Pro Phe Ile Leu Lys Ala Thr Gly Gly Pro Asn Leu Arg Gln 330
 Met Leu Pro Pro Pro 340
 345

<210> 11
 <211> 366
 <212> PRT
 <213> Arabidopsis thaliana

<400> 11
 Met Arg Pro Ile Gln Ser Pro Pro Gly Val Ser Val Pro Val Lys Ser
 1 5 10 15
 Arg Pro Arg Arg Arg Pro Asp Leu Thr Leu Pro Leu Pro Gln Arg Asp
 20 25 30
 Val Ser Leu Ala Val Pro Leu Pro Leu Pro Pro Thr Ser Gly Gly Ser
 35 40 45
 Gly Gly Ser Ser Gly Ser Ala Pro Ser Ser Gly Gly Ser Ala Ser Ser
 50 55 60
 Thr Asn Thr Asn Ser Ser Ile Glu Ala Lys Asn Tyr Ser Asp Leu Val
 65 70 75 80
 Arg Gly Asn Arg Ile Gly Ser Gly Ala Gly Gly Thr Val Tyr Lys Val
 85 90 95
 Ile His Arg Pro Ser Ser Arg Leu Tyr Ala Leu Lys Val Ile Tyr Gly
 100 105 110
 Asn His Glu Glu Thr Val Arg Arg Gln Ile Cys Arg Glu Ile Glu Ile
 115 120 125
 Leu Arg Asp Val Asn His Pro Asn Val Val Lys Cys His Glu Met Phe
 130 135 140
 Asp Gln Asn Gly Glu Ile Gln Val Leu Leu Glu Phe Met Asp Lys Gly
 145 150 155 160
 Ser Leu Glu Gly Ala His Val Trp Lys Glu Gln Gln Leu Ala Asp Leu
 165 170 175
 Ser Arg Gln Ile Leu Ser Gly Leu Ala Tyr Leu His Ser Arg His Ile
 180 185 190
 Val His Arg Asp Ile Lys Pro Ser Asn Leu Leu Ile Asn Ser Ala Lys
 195 200 205
 Asn Val Lys Ile Ala Asp Phe Gly Val Ser Arg Ile Leu Ala Gln Thr
 210 215 220
 Met Asp Pro Cys Asn Ser Ser Val Gly Thr Ile Ala Tyr Met Ser Pro
 225 230 235 240
 Glu Arg Ile Asn Thr Asp Leu Asn Gln Gly Lys Tyr Asp Gly Tyr Ala
 245 250 255
 Gly Asp Ile Trp Ser Leu Gly Val Ser Ile Leu Glu Phe Tyr Leu Gly
 260 265 270
 Arg Phe Pro Phe Pro Val Ser Arg Gln Gly Asp Trp Ala Ser Leu Met
 275 280 285
 Cys Ala Ile Cys Met Ser Gln Pro Pro Glu Ala Pro Ala Thr Ala Ser
 290 295 300
 Pro Glu Phe Arg His Phe Ile Ser Cys Cys Leu Gln Arg Glu Pro Gly
 305 310 315 320
 Lys Arg Arg Ser Ala Met Gln Leu Leu Gln His Pro Phe Ile Leu Arg
 325 330 335
 Ala Ser Pro Ser Gln Asn Arg Ser Pro Gln Asn Leu His Gln Leu Leu
 340 345 350
 Pro Pro Pro Arg Pro Leu Ser Ser Ser Ser Ser Pro Thr Thr
 355 360 365

<210> 12
 <211> 348
 <212> PRT
 <213> Arabidopsis thaliana

<400> 12
Met Lys Pro Ile Gln Ser Pro Ser Gly Val Ala Ser Pro Met Lys Asn
1 5 10 15
Arg Leu Arg Lys Arg Pro Asp Leu Ser Leu Pro Leu Pro His Arg Asp
20 25 30
Val Ala Leu Ala Val Pro Leu Pro Pro Ser Ser Ser Ser Ser
35 40 45
Ser Ala Pro Ala Ser Ser Ser Ala Ile Ser Thr Asn Ile Ser Ala Ala
50 55 60
Lys Ser Leu Ser Gln Leu Arg Val Asn Arg Ile Gly Ser Gly Ala
65 70 75 80
Gly Gly Thr Val Tyr Lys Val Ile His Thr Pro Thr Ser Arg Pro Phe
85 90 95
Ala Leu Lys Val Ile Tyr Gly Asn His Gln Asp Thr Val Arg Arg Gln
100 105 110
Ile Cys Arg Gln Ile Gln Ile Leu Arg Ser Val Asp His Pro Asn Val
115 120 125
Val Lys Cys His Asp Met Phe Asp His Asn Gly Gln Ile Gln Val Leu
130 135 140
Leu Gln Phe Met Asp Gln Gly Ser Leu Gln Gly Ala His Ile Trp Gln
145 150 155
Gln Gln Gln Leu Ala Asp Leu Ser Arg Gln Ile Leu Ser Gly Leu Ala
160 165 170 175
Tyr Leu His Arg Arg His Ile Val His Arg Asp Ile Lys Pro Ser Asn
180 185 190
Leu Leu Ile Asn Ser Ala Lys Asn Val Lys Ile Ala Asp Phe Gly Val
195 200 205
Ser Arg Ile Leu Ala Gln Thr Met Asp Pro Cys Asn Ser Ser Val Gly
210 215 220
Thr Ile Ala Tyr Met Ser Pro Gln Arg Ile Asn Thr Asp Leu Asn His
225 230 235
Gly Arg Tyr Asp Gly Tyr Ala Gly Asp Val Trp Ser Leu Gly Val Ser
240 245 250 255
Ile Leu Gln Phe Tyr Leu Gly Arg Phe Pro Phe Ala Val Ser Arg Gln
260 265 270 275
Gly Asp Trp Ala Ser Leu Met Cys Ala Ile Cys Met Ser Gln Pro Pro
280 285 290
Glu Ala Pro Ala Thr Ala Ser Gln Gln Phe Arg His Phe Val Ser Cys
295 300 305
Cys Leu Gln Ser Asp Pro Lys Arg Trp Ser Ala Gln Gln Leu Leu
310 315 320 325
Gln His Pro Phe Ile Leu Lys Ala Thr Gly Gly Pro Asn Leu Arg Gln
330 335 340
Met Leu Pro Pro Arg Pro Leu Pro Ser Ala Ser 345

<210> 13
<211> 2562
<212> DNA
<213> Arabidopsis thaliana

<400> 13
tcgtccatga tggacatata tgtgactgcg tattacaca caccgcacgt atgctattt 60
cacacgttag aagaagaatt caagaagtc ggttcattt gatatcttgt gagatcatca 120
atatgaccaat atcgttctat taatatagct ataattcata tgttggtcatg gtttcacata 180
ccatgctgac agtcgacgta cgtacaaaag tataaatagt atgaatctaa taacagcacc 240
aagatggaag ttcatcttct aatcaaaact atcataaagt ggtttcaaaa tagtggtttt 300

```

tctgatgaaa ctataactga gttataatca atccgaaatt atataactaa ttatatttgg 360
gaactagata aacgcaaaaa catgagcagt ttcttatttt ttttgtccag atttaaaatt 420
tgagtggtta aaatatacgg agtggttacac aatgaaaaca caagaagtca agaaccata 480
agttatttta attaataata ttgtatttaa agtgattatt aaaaaataat gtaaaaactg 540
attatttggt gacaaaaaac agttagttat agttaaatag tattgatgca tatatatact 600
atctcattat tttgggtatta ctccagtact acatctttta taaagacaaa gatagttagt 660
gtataattca aatcgaactc acagaagtca ataagcgcgt aaaaatacaa aaatatctgg 720
cagactttag caagggttgg tttccaacag aaatgggtcat ttcagaatca tttacatatc 780
catatatata gctcttaaat ggtatatatt gggtaatggg tattcgttta aataattttg 840
ttttctgtaa atttcaaata ttaatctgat cagtttatcc atgtgtgtat atttagtgta 900
ttatcatcaa tatatgacat agacagactt tcaagttggg gcaagagggg atgaaaattt 960
cttcccagtt gcaagagtaa gctgactagc attttttttt tatatataat ttatttctca 1020
aatgtgtttt attattggtt tgttgacttt aagtttttgc cttttatggg actgcaatca 1080
cccggtccaa cttttaatct ccctcgccta aaaaagaaag aaaaggctac cattatggac 1140
cgaaatattt aagaccataa tacaaaatta tacgaatatt ttctgtaact tatatatatg 1200
atcatttgac aaagcaaagt caatcaaata aacttcaaag aaattatgag cttataataa 1260
gtttgatagt gttaaatata atcaaatcat taaatttagt atttatttca tctcggtttc 1320
catttgatag atagataaat gaaagatagc atcgccaata atgaaaaaac tttatttgat 1380
ggcaataact tgttacatca tttctgtttt cttaatttca tgtcgaaata ttgccatgat 1440
tgtgttcaac ataactagtt ttgaggtaac aagttaaaaa tttgttatat tttttgaata 1500
tgttattcag ttgaaagtca ttttagatgta agtaaaaaaca aacataagaa gttaacatat 1560
caatatthaac acagcgaata atcattatta caaaaaaaag caaaaaaata gaagaagata 1620
ttatatattg gagaatcttt ctttagtctt agttgggaag attttgtgct atgggattaa 1680
aggtatccat cctatttcta tgatagaggc gtgggggtat tggaccaatc tataatatatt 1740
accacaaggc ttaagatgaa gtgataatac agtattatta ataccctccc aaattatttt 1800
taaataatta tcaaaagaag cttacgggat agatcactat tgcagcatta ttctataagt 1860
ttattttaatt tcagtggctc gttacgtgaa cacaaggtaa gctaataagac ttactgtccc 1920
cattaaacac atacataatt atacaagtat catgaaacta gtgacaaaac ctgcatacaa 1980
taaagaaatt accatgacga caaaagataa ttaaaaaaaa aactactata tgtcactact 2040
atgcatatgc atgtacaaat gccgctttta atatttaatt tagttaaaagc aatgatattt 2100
aaattctctc tacttcatat atattccaaa agacatatgg tcaaattcct ttttttagtt 2160
atatataatc atatatatcat attgttatat ttccaatatt taatagtaag atggactttc 2220
ctgaatgttg tgtatgattt ataatttgag atattttgtc ggagatggat atttgacaag 2280
ttaatgttac tttattaaaa ttttctaaac atttaggtac gaattgactt tttcaaaagt 2340
caacacaata aatttttaaaa gtttaatgac ttaacggggt cacatgggaa acgaaaacac 2400
cctaaaccac aaacaatcta atcttatttt cttctttata taaaccgctg tttcccaaaa 2460
ggcttgttct cgtcatatgt acttgtacac caaccaccca aaagagataa aagaggaaac 2520
aaaaactcga aaagagagag atatatgggt gaggtgggct at 2562

```

<210> 14

<211> 2796

<212> DNA

<213> Arabidopsis thaliana

<400> 14

```

ctgacagtga acttcattgt tcaagcgagg tgagtttctt atttttcttc ttctcttttc 60
aattaaaatt tcagggttta tgatctctag ggttttaggt ttattttctt aaactaaacc 120
ctaaattctt tttcttcttc tttttcttgg aatttccaga tgcactgcga acgctagagg 180
aggtgcgata tcgaagagga agcgttatac aggatcatgg gaccaaataa tatcgtatag 240
attcggttct tgttcttcta taccttatga gagtagaagt ttttctcttc aagagaaaaa 300
aaaaaaaaag aaaaaaaaaa aagaagagta gaaatttctc tgtgtttttt tttaccaaca 360
agacacaaat gaaactgggt caaaaggagt gtgtataatc tctgtggaga cataactaat 420
acgttgatga atttcaagaa tacttggatt atatagatta accctgactc ccttagatag 480
agatcgaaat cgggtggtga tttctgagac catacaagtt ctgatcagat tactactaag 540
taaactctta ttattagact gtttttaatg gatttttctg ctctaagat aacgtaatga 600
gatttttttt ttttttttgg atgtttgggt tgcagaccgg ggtaatgaga agtttggtta 660
gagcaaaagg cactaatctc acgtaagaaa acactttttt catcaaccat gtatataatc 720
atgtcgggtt acataaaccg tatcgtctat tcaagaattt agttttgtat aattataata 780

```

tttttttcag actacttttc aattaaagcat cttttttcttg gatcttttttt tcaacaaggg 840
 agagcttttaa cttttttgcat taactttat attttaaat atacaatgcat gcataccgac 900
 ttatatataat catatggtca atabgagac ttgtatgtat atatatgtc aacataagcat 960
 cttttccagat gaggtttcatg cacttttgt agaatattat ttgtatttat atatatgtc 1020
 gtttcaaaa gtttcaaaa aatccaatcc gttcaaaaat ttgtgaaact gttttgaaaga 1080
 ttgaaatggt ttgaggaagg atactcagtt ccaatctctg agcaggaatct gatattgactc 1140
 atctactcat aagaactttgc gagatagaac ggttcaaaaac cgtttccaag ggtttcataat 1200
 atabgagat aaagtgtgagtt ttgttgagcgt ttgttgagcgt ttgttgagcgt ttgttgagcgt 1260
 aacactaat agttatctcc ttgttgagcgt ttgttgagcgt ttgttgagcgt ttgttgagcgt 1320
 cgcagctatg tatatatttg gtttggttatc ttccaaaactc ttatatgtca aacaattttg 1380
 agaaagttat ttggaagttat aaaaagttat ttatatgtc agcactcctc 1440
 caaatatct ctttaaaact gtttgagcat aaaaagttat ttatatgtc agcactcctc 1500
 aatttaatt aacagccttt ttatcaaaa caaaaagttat ttatatgtc agcactcctc 1560
 aagttattat ggaagccttt ttatatgtc ttatatgtc ttatatgtc ttatatgtc 1620
 ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc 1680
 ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc 1740
 ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc 1800
 actattttgt tagttttgt tagttttgt tagttttgt tagttttgt tagttttgt 1860
 tagttttgt tagttttgt tagttttgt tagttttgt tagttttgt tagttttgt 1920
 aattttgt tagttttgt tagttttgt tagttttgt tagttttgt tagttttgt 1980
 aatatatatc tctttgttat aatatatatc tctttgttat aatatatatc tctttgttat 2040
 actttagaag ttgaaagaa aatatatatc tctttgttat aatatatatc tctttgttat 2100
 aaaaacatg ttgaaactg aatatatatc tctttgttat aatatatatc tctttgttat 2160
 cttcaaaaat ttgaaactg aatatatatc tctttgttat aatatatatc tctttgttat 2220
 ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc 2280
 ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc ttatatgtc 2340
 atagtttat actcaaaaag aatatatatc tctttgttat aatatatatc tctttgttat 2400
 tctttgttat actcaaaaag aatatatatc tctttgttat aatatatatc tctttgttat 2460
 aaaaatagat tttaagctc gtttttttat gcatatatgt ttatatgtc ttatatgtc 2520
 aatatatatc ttatatatat tttaagctc gtttttttat gcatatatgt ttatatgtc ttatatgtc 2580
 aatatatatc ttatatatat tttaagctc gtttttttat gcatatatgt ttatatgtc ttatatgtc 2640
 catgcaaat ttgaaatag tttaagctc gtttttttat gcatatatgt ttatatgtc ttatatgtc 2700
 gttttgttat gaaactcctc aagttttgt tttaagctc gtttttttat gcatatatgt ttatatgtc 2760
 gactatcaac atctatatg aaaaatggttt tttaagctc gtttttttat gcatatatgt ttatatgtc 2796